

PUBLISHED VERSION

Guo, Xuhong; Thomas, Anthony William; Williams, Anthony Gordon
[Bethe-Salpeter equation for heavy baryons \$\omega_Q^*\$ in the diquark picture](#) Physical Review D,
1999; 59(11):116007

©1999 American Physical Society

<http://link.aps.org/doi/10.1103/PhysRevD.59.116007>

PERMISSIONS

<http://publish.aps.org/authors/transfer-of-copyright-agreement>

“The author(s), and in the case of a Work Made For Hire, as defined in the U.S. Copyright Act, 17 U.S.C.

§101, the employer named [below], shall have the following rights (the “Author Rights”):

[...]

3. The right to use all or part of the Article, including the APS-prepared version without revision or modification, on the author(s)' web home page or employer's website and to make copies of all or part of the Article, including the APS-prepared version without revision or modification, for the author(s)' and/or the employer's use for educational or research purposes.”

17th April 2013

<http://hdl.handle.net/2440/10895>

Bethe-Salpeter equation for heavy baryons $\omega_Q^{(*)}$ in the diquark picture

X.-H. Guo*

*Department of Physics and Mathematical Physics, and Special Research Center for the Subatomic Structure of Matter;
University of Adelaide, South Australia 5005, Australia;
and Institute of High Energy Physics, Academia Sinica, Beijing 100039, China*

A. W. Thomas[†] and A. G. Williams[‡]

*Department of Physics and Mathematical Physics, and Special Research Center for the Subatomic Structure of Matter,
University of Adelaide, South Australia 5005, Australia*

(Received 13 May 1998; revised manuscript received 2 December 1998; published 27 April 1999)

In the heavy quark limit, the heavy baryons $\omega_Q^{(*)}$ (ω stands for Σ , Ξ or Ω and $Q=b$ or c) are regarded as composed of a heavy quark and an axial vector, light diquark with good spin and isospin quantum numbers. Based on this diquark picture we establish the Bethe-Salpeter (BS) equation for $\omega_Q^{(*)}$ in the limit where the heavy quark has infinite mass, $m_Q \rightarrow \infty$. It is found that in this limit there are three components in the BS wave function for $\omega_Q^{(*)}$. Assuming the kernel to consist of a scalar confinement term and a one-gluon-exchange term we derive three coupled integral equations for the three BS scalar functions in the covariant instantaneous approximation. Numerical solutions for the three BS scalar functions are presented, including a discussion of their dependence on the various input parameters. These solutions are applied to calculate the Isgur-Wise functions $\xi(\omega)$ and $\zeta(\omega)$ for the weak transitions $\Omega_b^{(*)} \rightarrow \Omega_c^{(*)}$. Using these we give predictions for the Cabibbo-allowed nonleptonic decay widths and up-down asymmetries for $\Omega_b \rightarrow \Omega_c^{(*)}$ plus a pseudoscalar or vector meson. [S0556-2821(99)07909-6]

PACS number(s): 11.10.St, 12.39.Hg, 14.20.Lq, 14.20.Mr

I. INTRODUCTION

The physics of heavy hadrons has been a subject of intense interest in recent years. One reason for this is that more and more experimental data are being accumulated. Another reason is the discovery of the new flavor and spin symmetry in QCD, $SU(2)_f \times SU(2)_s$, in the heavy quark limit and the establishment of heavy quark effective theory (HQET) [1]. However, in comparison with the heavy meson case, heavy baryons have been studied much less, both experimentally and theoretically.

On the other hand, the experimental situation with heavy baryons has been improving recently, with more measurements becoming available. For instance, OPAL has measured some physical quantities for Λ_b , such as its lifetime and the production branching ratio for the inclusive semileptonic decay $\Lambda_b \rightarrow \Lambda l^- \bar{\nu}_X$ [2]. Furthermore, measurements of the nonleptonic decay of Λ_b have also been made, through the well-known process $\Lambda_b \rightarrow \Lambda J/\psi$. The discrepancy between the measurements made by UA1 [3] and those by the Collider Detector at Fermilab (CDF) [4] and CERN e^+e^- collider LEP Collaborations [5] appears to have been settled by the new measurement from CDF [6]. However, compared with D and B mesons, the data for heavy baryons are still very limited. In addition to Λ_b , there have been few data on other bottom baryons [7], although we expect more data to appear in the near future. Clearly the time is right for serious

theoretical studies of heavy baryon properties to begin.

Theoretically, HQET can simplify the physical processes involving heavy quarks, since with the aid of HQET the number of independent form factors is reduced. For instance, in leading order of the $1/m_Q$ expansion only one form factor (the Isgur-Wise function) remains for the $\Lambda_b \rightarrow \Lambda_c$ transition, while for $\omega_b^{(*)} \rightarrow \omega_c^{(*)}$ (we follow the notation of Ref. [8]; ω could be Ξ , Σ or Ω and $Q=b$ or c) there are two independent Isgur-Wise functions. (Note that $\omega_Q^{(*)}$ is a notation implying either ω_Q or ω_Q^* .) The behavior of these functions depends on the nonperturbative effects of QCD which control the dynamics inside a heavy hadron. Hence some nonperturbative QCD model has to be adopted from which these Isgur-Wise functions can be obtained. In previous work [9], we established the Bethe-Salpeter (BS) equation for Λ_Q , which is assumed to be composed of a heavy quark, Q , and a scalar diquark. Some theoretical predictions for $\Lambda_b \rightarrow \Lambda_c$ were also obtained. It is the purpose of the present paper to generalize such an approach to the heavy baryons, $\omega_Q^{(*)}$, and consequently give some phenomenological predictions for the weak decays of such baryons.

When the quark mass is very heavy compared with the QCD scale, Λ_{QCD} , the light degrees of freedom in a heavy baryon, Λ_Q ($Q=b$ or c), become blind to the flavor and spin quantum numbers of the heavy quark because of the $SU(2)_f \times SU(2)_s$ symmetry. Therefore, the light degrees of freedom have good quantum numbers, including angular momentum and isospin. These quantum numbers can be used to classify heavy baryons. For example, the light degrees of freedom of Λ_Q have zero angular momentum and isospin. For $\Sigma_Q^{(*)}$, the angular momentum and parity J^P of the light degrees of freedom are 1^+ , and the isospin is also 1 in order to guarantee that the total wave function of the hadron is antisymmetric. Hence it is natural to consider the heavy

*Email address: xhguo@physics.adelaide.edu.au

†Email address: athomas@physics.adelaide.edu.au

‡Email address: awilliam@physics.adelaide.edu.au

baryon to be composed of a heavy quark and a light diquark. This is our underlying assumption.

Based on the picture of the composition of the heavy baryon which we have just presented, the three body system is simplified to a two body system. We establish the BS equation for the heavy baryons, $\omega_Q^{(*)}$, in this picture. The heavy quark symmetry can be used to simplify the form of the BS wave function greatly. It can be shown that in the limit $m_Q \rightarrow \infty$ there are three components in the BS wave function, and hence we have three corresponding scalar functions. We solve the BS equation numerically by assuming that the kernel contains a scalar confinement term and a one-gluon-exchange term. The explicit dependence of the BS wave function on the parameters of the model will be discussed. Furthermore, we calculate the Isgur-Wise functions in terms of the BS wave functions and give theoretical predictions for the Cabibbo-allowed two body nonleptonic decays $\Omega_b \rightarrow \Omega_c^{(*)}$ plus a pseudoscalar or vector meson.

The light degrees of freedom of $\omega_Q^{(*)}$ belong to a 6 representation of flavor SU(3). Taking $Q=b$ as an example, $\omega_b^{(*)}$ includes $\Sigma_b^{(*)+,0,-}$, $\Xi_b^{(*)0,-}$ and $\Omega_b^{(*)-}$. The total spins of ω_Q and ω_Q^* are $\frac{1}{2}$ and $\frac{3}{2}$ respectively. There is no strange quark in $\Sigma_Q^{(*)}$ while there is one strange quark in $\Xi_Q^{(*)}$ and two in $\Omega_Q^{(*)}$ respectively.

The remainder of this paper is organized as follows. In Sec. II we establish the BS equation for the heavy quark and axial vector light diquark system and discuss the form of the kernel. In Sec. III we derive explicitly the coupled integral equations for the BS scalar wave functions. In Sec. IV we discuss the normalization condition of the BS wave function by exploiting the normalization of the Isgur-Wise function at the zero recoil point. The numerical solutions of the BS equation and their dependence on the parameters in our model are presented in Sec. V. In Sec. VI we calculate the Isgur-Wise functions and give predictions for the decay widths and up-down asymmetry parameters for $\Omega_b \rightarrow \Omega_c^{(*)}$ plus a pseudoscalar or vector meson. Finally, Sec. VI contains a summary and discussions.

II. BS EQUATION FOR $\omega_Q^{(*)}$

As discussed in Sec. I, $\omega_Q^{(*)}$ is regarded as a bound state of a heavy quark, ψ_Q , and a light axial vector diquark, A_μ . In the following, u_Q denotes the Dirac spinor of ψ_Q and η_μ represents the polarization vector of A_μ . Then $B_\mu \equiv u_Q \eta_\mu$ can be decomposed into spin- $\frac{1}{2}$ and spin- $\frac{3}{2}$ states [10,11] which represent ω_Q and ω_Q^* respectively. These two states are degenerate in the heavy quark limit. Using the notation of Refs. [11,12] this doublet is described by $B_\mu^m(v)$, where $m=1,2$ correspond to ω_Q and ω_Q^* respectively, v_μ is the velocity of the heavy baryon and $B_\mu = B_\mu^1 + B_\mu^2$ [10]. Explicitly, we can write

$$\begin{aligned} B_\mu^1(v) &= \frac{1}{\sqrt{3}}(\gamma_\mu + v_\mu)\gamma_5 u(v), \\ B_\mu^2(v) &= u_\mu(v), \end{aligned} \quad (1)$$

where $u(v)$ is the Dirac spinor and $u_\mu(v)$ is the Rarita-Schwinger vector spinor. $B_\mu^m(v)$ satisfies the following conditions:

$$\not{v} B_\mu^m(v) = B_\mu^m(v), \quad v^\mu B_\mu^m(v) = 0, \quad \gamma^\mu B_\mu^2(v) = 0. \quad (2)$$

The above constraints for $m=1$ can be seen from $\not{v} u(v) = u(v)$ while for $m=2$, they are the properties of a spin- $\frac{3}{2}$, Rarita-Schwinger vector spinor.

Under a Lorentz transformation Λ [10],

$$B_\mu \rightarrow \Lambda_\mu^\nu D(\Lambda) B_\nu, \quad (3)$$

where $D(\Lambda)$ is the spinorial representation of Λ . Under the heavy quark spin transformation [8],

$$B_\mu \rightarrow -\gamma_5 \not{v} \not{e} B_\mu, \quad (4)$$

where $e = e_1, e_2, e_3$ are three mutually orthogonal four-dimensional unit vectors which are also orthogonal to v (i.e., $e_i \cdot v = 0$) and $e_i^2 = -1$ ($i=1,2,3$). The three unit vectors are associated with the heavy quark spin operators which are generators of the $SU(2)_s$ symmetry.

Since $\omega_Q^{(*)}$ is composed of ψ_Q and A_μ , we can define the BS wave function of $\omega_Q^{(*)}$ by

$$\chi_\mu(x_1, x_2, P) = \langle 0 | T \psi_Q(x_1) A_\mu(x_2) | \omega_Q^{(*)}(P) \rangle, \quad (5)$$

where $P = m_{\omega_Q^{(*)}} v$ is the total momentum of $\omega_Q^{(*)}$ and v is its velocity. Let m_Q and m_D be the masses of the heavy quark and the light diquark in the baryon. Let us define $\lambda_1 \equiv m_Q / (m_Q + m_D)$ and $\lambda_2 \equiv m_D / (m_Q + m_D)$ and let p be the relative momentum of the two constituents. The BS wave function in momentum space is defined as

$$\chi_\mu^m(x_1, x_2, P) = e^{iP \cdot X} \int \frac{d^4 p}{(2\pi)^4} e^{ip \cdot x} \chi_{Pm}^m(p), \quad (6)$$

where $X \equiv \lambda_1 x_1 + \lambda_2 x_2$ is the coordinate of the center of mass and $x \equiv x_1 - x_2$. The momentum of the heavy quark is $p_1 = \lambda_1 P + p$ and that of the diquark is $p_2 = -\lambda_2 P + p$.

It can be shown that $\chi_{Pm}^\mu(p)$ satisfies the following BS equation [13]

$$\begin{aligned} \chi_{Pm}^\mu(p) &= S_F(\lambda_1 P + p) \int \frac{d^4 q}{(2\pi)^4} G_{\rho\nu}(P, p, q) \\ &\quad \times \chi_{Pm}^\nu(q) S_D^{\mu\rho}(-\lambda_2 P + p), \end{aligned} \quad (7)$$

where $G_{\rho\nu}(P, p, q)$ is the kernel, which is defined as the sum of all the two particle irreducible diagrams with respect to the heavy quark and the light diquark. For convenience, in the following we use the variables

$$p_l \equiv v \cdot p - \lambda_2 m_{\omega_Q^{(*)}}, \quad p_i \equiv p - (v \cdot p)v. \quad (8)$$

Then in the leading order of the $1/m_Q$ expansion we have

$$S_F(\lambda_1 P + p) = \frac{i(1 + \not{v})}{2(p_l + E_0 + m_D + i\epsilon)}, \quad (9)$$

where E_0 is the binding energy. From Eqs. (7) and (9) it follows that $\chi_{Pm}^\mu(p)$ satisfies the following equation:

$$\not{v}\chi_{Pm}^\mu(p) = \chi_{Pm}^\mu(p). \quad (10)$$

The propagator of the light axial vector diquark has the form

$$S_{D\mu\nu}(p_2) = -i \frac{g_{\mu\nu} - p_{2\mu}p_{2\nu}/m_D^2}{p_2^2 - m_D^2 + i\epsilon}. \quad (11)$$

In the limit $m_Q \rightarrow \infty$ we have $p_2 = -m_D v + p$ and hence we have

$$\begin{aligned} S_{D\mu\nu}(p_2) \\ = -i \frac{g_{\mu\nu} - v_\mu v_\nu - p_\mu p_\nu / m_D^2 + (v_\mu p_\nu + v_\nu p_\mu) / m_D}{p_1^2 - W_p^2 + i\epsilon}, \end{aligned} \quad (12)$$

where $W_p \equiv \sqrt{p_1^2 + m_D^2}$. The corrections to Eqs. (9) and (12) are from $O(1/m_Q)$ terms.

Now we discuss the form of the BS wave function $\chi_{Pm}^\mu(p)$. In the heavy quark limit, as a result of the $SU(2)_s \times SU(2)_f$ symmetry, the internal dynamics of the heavy baryon, $\omega_Q^{(*)}$, is determined by the light degrees of freedom and the flavor and spin direction of the heavy quark, Q , is irrelevant. Consequently we have [10,12]

$$\chi_P^\mu(p) = u_Q(v) \eta_\nu \zeta^{\mu\nu}(v, p) \quad (13)$$

and

$$\langle 0 | A_\mu | \text{light}, 1^+ \rangle = \eta_\nu \zeta^{\mu\nu}(v, p). \quad (14)$$

Since $v \cdot \eta = 0$ [10], the tensor $\zeta^{\mu\nu}(v, p)$ can be expanded as

$$\zeta^{\mu\nu}(v, p) = A_1 g^{\mu\nu} + A_2 v^\mu p^\nu + A_3 p^\mu p^\nu, \quad (15)$$

where $A_i (i=1,2,3)$ are Lorentz scalar functions. After expressing $u_Q \eta_\mu$ in Eq. (13) in terms of $B_\mu^m(v) (m=1,2)$, we have the following form for the BS wave function:

$$\chi_{Pm}^\mu = A_1 B_m^\mu(v) + A_2 v^\mu p_\nu B_m^\nu(v) + A_3 p^\mu p_\nu B_m^\nu(v). \quad (16)$$

Therefore, we have three components in the BS wave function, $\chi_{Pm}^\mu(p)$, and they correspond to three scalar BS functions $A_i (i=1,2,3)$. This is consistent with our diquark picture for $\omega_Q^{(*)}$. In the heavy quark limit, the dynamics inside the heavy baryon is controlled by the configuration of the light degrees of freedom. Since the light diquark is a 1^+ object, it has three different configurations. Consequently there are three components in the BS wave function which describe the dynamics in the heavy baryon $\omega_Q^{(*)}$.

In fact, we can derive the form of $\chi_{Pm}^\mu(p)$ in another way. We may first write out all the possible terms which have the same behavior as $\chi_{Pm}^\mu(p)$ under Lorentz transformations. Then by applying the condition, Eq. (10), and ensuring the proper behavior under the heavy quark spin transformation, Eq. (4), we obtain the same result as given in Eq. (16).




FIG. 1. The vertex of two axial vector diquarks and a gluon.

Considering $p^\mu = v \cdot p v^\mu + p_1^\mu$, and using the constraint $v^\mu B_\mu^m(v) = 0$, it will be convenient to define

$$A = A_1, \quad C = A_2 + v \cdot p A_3, \quad D = A_3,$$

which results in the following expression for the BS wave function:

$$\chi_{Pm}^\mu = A B_m^\mu(v) + C v^\mu p_{1\nu} B_m^\nu(v) + D p_1^\mu p_{1\nu} B_m^\nu(v). \quad (17)$$

A, C and D in Eq. (17) are functions of p_1 and p_1^2 . Their behavior is controlled by nonperturbative QCD. Our aim is to obtain explicit forms for them with some QCD-motivated model for the form of the BS kernel.

Motivated by the success of the potential model [14], scalar confinement and one-gluon-exchange terms were used in the kernel when studying Λ_Q in Ref. [9]. This form was also used in the heavy meson case in Ref. [15]. In the present work we will also adopt this form of the kernel

$$iG^{\rho\nu} = g^{\rho\nu} I \otimes IV_1 + v_\mu \otimes \Gamma^{\mu\rho\nu} V_2, \quad (18)$$

where $\Gamma^{\mu\rho\nu}$ is the vertex of a gluon with two axial vector diquarks. This vertex should reflect the internal structure of the diquark. In this work, we use the model proposed in Ref. [16] where this vertex has the following form (see Fig. 1):

$$-i \frac{\lambda^a}{2} g_s \Gamma^{\mu\rho\nu} F_V(Q^2), \quad (19)$$

with

$$\Gamma^{\mu\rho\nu} = (p_2 + p_2')^\mu g^{\nu\rho} - (p_2^\nu g^{\mu\rho} + p_2'^\rho g^{\mu\nu}).$$

In Eq. (19), g_s is the strong interaction coupling constant and $F_V(Q^2)$ is introduced to describe the internal structure of the axial vector diquark. The form factor, $F_V(Q^2)$, depends on nonperturbative QCD interactions and will be determined phenomenologically, by comparison with experiment.

As discussed in Ref. [9], when we consider the vertex of two heavy quarks with a gluon, the momenta of the two heavy quarks are $p_1 = \lambda_1 m_{\Lambda_Q} v + p$ and $p_1' = \lambda_1 m_{\Lambda_Q} v + q$ respectively, where p and q are relative momenta and of the order Λ_{QCD} . In the heavy quark limit the heavy quark is almost on-shell and moves with constant velocity. It can be shown that $p_1 = q_1$ at this vertex when the heavy quark is

exactly on-shell. This is the so-called covariant instantaneous approximation [9,15]. With this approximation, V_1 and V_2 in $G^{\rho\nu}(P,p,q)$ are replaced by

$$\tilde{V}_i \equiv V_i|_{p_l=q_l} \quad (i=1,2). \quad (20)$$

III. COUPLED INTEGRAL EQUATIONS FOR THREE BS SCALAR WAVE FUNCTIONS

In this section we will derive explicitly three coupled integral equations for the BS scalar wave functions. Substituting Eqs. (9) and (12) into Eq. (7) and considering the form of the kernel in Eq. (18) and the property in Eq. (10), we obtain the following form for the BS equation

$$\chi_{Pm}^\mu(p) = \frac{-i}{(p_l + E_0 + m_D + i\epsilon)(p_l^2 - W_p^2 + i\epsilon)} M_m^\mu, \quad (21)$$

where

$$M_{m\mu} \equiv i \left[g_{\mu\rho} - v_\mu v_\rho + \frac{(v_\mu p_\rho + p_\mu v_\rho)}{m_D} - \frac{p_\mu p_\rho}{m_D^2} \right] \times \int \frac{d^4 q}{(2\pi)^4} [G^{\rho\nu}(P,p,q) \chi_{Pm\nu}(q)]|_{p_l=q_l}, \quad (22)$$

and we have made explicit use of the covariant instantaneous approximation.

Substituting Eqs. (17), (18) and (19) into Eq. (22) and again using the covariant instantaneous approximation we have

$$\begin{aligned} M_m^\mu = & B_m^\mu(v) \int \frac{d^4 q}{(2\pi)^4} A(\tilde{V}_1 + 2p_l \tilde{V}_2) + \frac{1}{m_D} v^\mu \int \frac{d^4 q}{(2\pi)^4} \{ p_t \cdot B_m(v) A(\tilde{V}_1 + 2p_l \tilde{V}_2) + (p_l + m_D) [-A p_t \cdot B_m(v) \tilde{V}_2 \\ & + q_t \cdot B_m(v) (C \tilde{V}_1 - D p_t \cdot q_t \tilde{V}_2)] + p_t \cdot q_t q_t \cdot B_m(v) [-C \tilde{V}_2 + D(\tilde{V}_1 + 2p_l \tilde{V}_2)] \} \\ & - \frac{1}{m_D^2} p^\mu \int \frac{d^4 q}{(2\pi)^4} \{ p_t \cdot B_m(v) A(\tilde{V}_1 + 2p_l \tilde{V}_2) + p_l [-A p_t \cdot B_m(v) \tilde{V}_2 + q_t \cdot B_m(v) (C \tilde{V}_1 - D p_t \cdot q_t \tilde{V}_2)] \\ & + p_t \cdot q_t q_t \cdot B_m(v) [-C \tilde{V}_2 + D(\tilde{V}_1 + 2p_l \tilde{V}_2)] \} + \int \frac{d^4 q}{(2\pi)^4} q_t^\mu q_t \cdot B_m(v) [-C \tilde{V}_2 + D(\tilde{V}_1 + 2p_l \tilde{V}_2)]. \end{aligned} \quad (23)$$

We notice that in Eq. (23) there are terms of the form $\int [d^4 q / (2\pi)^4] q_t^\mu f$ and $\int [d^4 q / (2\pi)^4] q_t^\mu q_t^\nu f$, where f is some function of p^2, q^2 and $p \cdot q$. On the grounds of Lorentz invariance, in general we have

$$\int \frac{d^4 q}{(2\pi)^4} q_t^\mu f = f_1 v^\mu + f_2 p_t^\mu \quad (24)$$

and

$$\int \frac{d^4 q}{(2\pi)^4} q_t^\mu q_t^\nu f = g_1 g^{\mu\nu} + g_2 v^\mu v^\nu + g_3 v^\mu p_t^\nu + g_4 v^\nu p_t^\mu + g_5 p_t^\mu p_t^\nu. \quad (25)$$

From Eq. (24) only the f_2 term can contribute, while from Eq. (25) only the g_1, g_3 and g_5 terms can contribute, since $v_\nu B_m^\nu(v) = 0$. It can be easily shown that

$$\begin{aligned} f_2 &= \int \frac{d^4 q}{(2\pi)^4} \frac{p_t \cdot q_t}{p_t^2} f, \\ g_1 &= \int \frac{d^4 q}{(2\pi)^4} \frac{(p_t \cdot q_t)^2 - p_t^2 q_t^2}{2p_t^2} f, \\ g_3 &= 0, \\ g_5 &= \int \frac{d^4 q}{(2\pi)^4} \frac{3(p_t \cdot q_t)^2 - p_t^2 q_t^2}{2p_t^4} f. \end{aligned} \quad (26)$$

With the aid of Eqs. (24), (25) and (26) we can express M_m^μ in Eq. (23) in terms of $B_m^\mu(v)$, $v^\mu p_{t\nu} B_m^\nu(v)$ and $p_t^\mu p_{t\nu} B_m^\nu(v)$. Let us define

$$\tilde{A}(p_t^2) = \int \frac{dq_l}{2\pi} A(p_l, p_t^2), \quad \tilde{C}(p_t^2) = \int \frac{dq_l}{2\pi} C(p_l, p_t^2), \quad \tilde{D}(p_t^2) = \int \frac{dq_l}{2\pi} D(p_l, p_t^2), \quad (27)$$

where \tilde{A} , \tilde{C} and \tilde{D} are functions of p_i^2 only. Then one obtains the expression

$$\begin{aligned}
M_m^\mu = & B_m^\mu(v) \int \frac{d^3 q_t}{(2\pi)^3} \left\{ \tilde{A}(\tilde{V}_1 + 2p_l \tilde{V}_2) - \tilde{C} \frac{(p_t \cdot q_t)^2 - p_t^2 q_t^2}{2p_t^2} \tilde{V}_2 + \tilde{D} \frac{(p_t \cdot q_t)^2 - p_t^2 q_t^2}{2p_t^2} (\tilde{V}_1 + 2p_l \tilde{V}_2) \right\} \\
& + \frac{1}{m_D^2} v^\mu p_t \cdot B_m(v) \int \frac{d^3 q_t}{(2\pi)^3} \left\{ -\tilde{A}[p_l \tilde{V}_1 + (p_l^2 + m_D^2) \tilde{V}_2] - \tilde{C} \left[(p_l^2 - m_D^2) \frac{p_t \cdot q_t}{p_t^2} \tilde{V}_1 + p_l \frac{(p_t \cdot q_t)^2}{p_t^2} \tilde{V}_2 \right] \right. \\
& + \tilde{D} \frac{(p_t \cdot q_t)^2}{p_t^2} [p_l \tilde{V}_1 + (p_l^2 + m_D^2) \tilde{V}_2] \left. \right\} - \frac{1}{m_D^2} p_t^\mu p_t \cdot B_m(v) \int \frac{d^3 q_t}{(2\pi)^3} \left\{ \tilde{A}(\tilde{V}_1 + p_l \tilde{V}_2) \right. \\
& + \tilde{C} \left[p_l \frac{p_t \cdot q_t}{p_t^2} \tilde{V}_1 + \frac{m_D^2(3(p_t \cdot q_t)^2 - p_t^2 q_t^2) + 2p_t^2(p_t \cdot q_t)^2}{2p_t^4} \tilde{V}_2 \right] \\
& \left. + \tilde{D} \left[-\frac{m_D^2(3(p_t \cdot q_t)^2 - p_t^2 q_t^2) + 2p_t^2(p_t \cdot q_t)^2}{2p_t^4} (\tilde{V}_1 + 2p_l \tilde{V}_2) + p_l \frac{(p_t \cdot q_t)^2}{p_t^2} \tilde{V}_2 \right] \right\}. \tag{28}
\end{aligned}$$

In Eq. (21) there are poles in p_l at $W_p - i\epsilon$, $-W_p + i\epsilon$ and $-E_0 - m_D - i\epsilon$. By choosing the appropriate contour, we integrate over p_l on both sides of Eq. (21) and obtain the following three coupled integral equations for \tilde{A} , \tilde{C} and \tilde{D} :

$$\begin{aligned}
\tilde{A}(p_i^2) = & \frac{-1}{2W_p(E_0 + m_D - W_p)} \int \frac{d^3 q_t}{(2\pi)^3} \left\{ \tilde{A}(q_t^2)(\tilde{V}_1 - 2W_p \tilde{V}_2) - \tilde{C}(q_t^2) \frac{(p_t \cdot q_t)^2 - p_t^2 q_t^2}{2p_t^2} \tilde{V}_2 \right. \\
& \left. + \tilde{D}(q_t^2) \frac{(p_t \cdot q_t)^2 - p_t^2 q_t^2}{2p_t^2} (\tilde{V}_1 - 2W_p \tilde{V}_2) \right\}, \tag{29}
\end{aligned}$$

$$\begin{aligned}
\tilde{C}(p_i^2) = & \frac{-1}{2m_D^2 W_p(E_0 + m_D - W_p)} \int \frac{d^3 q_t}{(2\pi)^3} \left\{ \tilde{A}(q_t^2) \{W_p \tilde{V}_1 - [(E_0 + m_D)W_p + m_D^2] \tilde{V}_2\} \right. \\
& + \tilde{C}(q_t^2) \left[-\frac{p_t \cdot q_t}{p_t^2} [(E_0 + m_D)W_p - m_D^2] \tilde{V}_1 + W_p \frac{(p_t \cdot q_t)^2}{p_t^2} \tilde{V}_2 \right] \\
& \left. + \tilde{D}(q_t^2) \frac{(p_t \cdot q_t)^2}{p_t^2} \{-W_p \tilde{V}_1 + [(E_0 + m_D)W_p + m_D^2] \tilde{V}_2\} \right\}, \tag{30}
\end{aligned}$$

$$\begin{aligned}
\tilde{D}(p_i^2) = & \frac{1}{2m_D^2 W_p(E_0 + m_D - W_p)} \int \frac{d^3 q_t}{(2\pi)^3} \left\{ \tilde{A}(q_t^2)(\tilde{V}_1 - W_p \tilde{V}_2) \right. \\
& + \tilde{C}(q_t^2) \left[-\frac{p_t \cdot q_t}{p_t^2} W_p \tilde{V}_1 + \frac{m_D^2[3(p_t \cdot q_t)^2 - p_t^2 q_t^2] + 2p_t^2(p_t \cdot q_t)^2}{2p_t^4} \tilde{V}_2 \right] \\
& \left. - \tilde{D}(q_t^2) \left[\frac{m_D^2[3(p_t \cdot q_t)^2 - p_t^2 q_t^2] + 2p_t^2(p_t \cdot q_t)^2}{2p_t^4} (\tilde{V}_1 - 2W_p \tilde{V}_2) + \frac{(p_t \cdot q_t)^2}{p_t^2} W_p \tilde{V}_2 \right] \right\}. \tag{31}
\end{aligned}$$

If one knows the form for the kernel, \tilde{V}_1 and \tilde{V}_2 , then $\tilde{A}(p_i^2)$, $\tilde{C}(p_i^2)$ and $\tilde{D}(p_i^2)$ can be obtained from Eqs. (29), (30) and (31). Consequently from Eqs. (17), (21) and (28) we find the following expressions for $A(p_l, p_i^2)$, $C(p_l, p_i^2)$ and $D(p_l, p_i^2)$:

$$\begin{aligned}
A(p_l, p_i^2) = & \frac{-i}{(p_l + E_0 + m_D + i\epsilon)(p_l^2 - W_p^2 + i\epsilon)} \int \frac{d^3 q_t}{(2\pi)^3} \left\{ \tilde{A}(q_t^2)(\tilde{V}_1 + 2p_l \tilde{V}_2) - \tilde{C}(q_t^2) \frac{(p_t \cdot q_t)^2 - p_t^2 q_t^2}{2p_t^2} \tilde{V}_2 \right. \\
& \left. + \tilde{D}(q_t^2) \frac{(p_t \cdot q_t)^2 - p_t^2 q_t^2}{2p_t^2} (\tilde{V}_1 + 2p_l \tilde{V}_2) \right\}, \tag{32}
\end{aligned}$$

$$C(p_l, p_t^2) = \frac{-i}{m_D^2(p_l + E_0 + m_D + i\epsilon)(p_l^2 - W_p^2 + i\epsilon)} \int \frac{d^3 q_t}{(2\pi)^3} \left\{ -\tilde{A}(q_t^2)[p_l \tilde{V}_1 + (p_l^2 + m_D^2) \tilde{V}_2] \right. \\ \left. - \tilde{C}(q_t^2) \left[(p_l^2 - m_D^2) \frac{p_l \cdot q_t}{p_t^2} \tilde{V}_1 + p_l \frac{(p_l \cdot q_t)^2}{p_t^2} \tilde{V}_2 \right] + \tilde{D}(q_t^2) \frac{(p_l \cdot q_t)^2}{p_t^2} [p_l \tilde{V}_1 + (p_l^2 + m_D^2) \tilde{V}_2] \right\}, \quad (33)$$

$$D(p_l, p_t^2) = \frac{i}{m_D^2(p_l + E_0 + m_D + i\epsilon)(p_l^2 - W_p^2 + i\epsilon)} \int \frac{d^3 q_t}{(2\pi)^3} \left\{ \tilde{A}(q_t^2)(\tilde{V}_1 + p_l \tilde{V}_2) \right. \\ \left. + \tilde{C}(q_t^2) \left[\frac{p_l \cdot q_t}{p_t^2} p_l \tilde{V}_1 + \frac{m_D^2[3(p_l \cdot q_t)^2 - p_t^2 q_t^2] + 2p_t^2(p_l \cdot q_t)^2}{2p_t^4} \tilde{V}_2 \right] \right. \\ \left. + \tilde{D}(q_t^2) \left[-\frac{m_D^2[3(p_l \cdot q_t)^2 - p_t^2 q_t^2] + 2p_t^2(p_l \cdot q_t)^2}{2p_t^4} (\tilde{V}_1 + 2p_l \tilde{V}_2) + \frac{(p_l \cdot q_t)^2}{p_t^2} p_l \tilde{V}_2 \right] \right\}. \quad (34)$$

A model kernel, specified in terms of \tilde{V}_1 and \tilde{V}_2 , for the BS equation in the scalar light diquark case was given in Ref. [9]. In the present axial vector diquark case, a model vertex for a gluon with two 1^+ diquarks is given in Eq. (19), where $F_V(Q^2)$ describes the internal structure of the light diquark. Following Ref. [16] we take the form for $F_V(Q^2)$ as

$$F_V(Q^2) = \frac{\alpha_s^{(\text{eff})} Q_1^2}{Q^2 + Q_1^2}, \quad (35)$$

where Q_1^2 is a parameter which freezes $F_V(Q^2)$ when Q^2 is very small. In the high energy region the form factor is proportional to $1/Q^2$, which is consistent with perturbative QCD calculations [17]. By analyzing the electromagnetic form factor for the proton it was found that selecting $Q_1^2 = 3.2 \text{ GeV}^2$ can lead to results consistent with the experimental data [16]. Note that in Eq. (35) we do not consider the difference between longitudinal and transverse polarization states. The reason is that we are considering the bound state with a binding energy of order Λ_{QCD} ; so this difference should be small [see Ref. [16], where this difference is a factor $Q_2^2/(Q^2 + Q_2^2)$ with Q_2^2 approximately 15 GeV^2 ; so this factor is close to 1 in our discussion].

Based on the above discussion, the kernel for the BS equation in the baryon case is taken to have the following form:

$$\tilde{V}_1 = \frac{8\pi\kappa}{[(p_t - q_t)^2 + \mu^2]^2} - (2\pi)^3 \delta^3(p_t - q_t) \\ \times \int \frac{d^3 k}{(2\pi)^3} \frac{8\pi\kappa}{(k^2 + \mu^2)^2}, \\ \tilde{V}_2 = -\frac{16\pi}{3} \frac{\alpha_s^{(\text{eff})2} Q_1^2}{[(p_t - q_t)^2 + \mu^2][(p_t - q_t)^2 + Q_1^2]}, \quad (36)$$

where κ and $\alpha_s^{(\text{eff})}$ are coupling parameters related to scalar confinement and the one-gluon-exchange diagram respectively. The second term of \tilde{V}_1 in Eq. (36) is the counter-term which removes the infra-red divergence arising from the linear confinement in the integral equation. The parameter μ is introduced to avoid the infra-red divergence in numerical calculations. The limit $\mu \rightarrow 0$ is taken in the end. Besides Q_1^2 , there are two parameters, κ and $\alpha_s^{(\text{eff})}$, in the kernel. However, they should be related to each other when we solve the coupled integral equations (29), (30) and (31). This will be discussed in detail in Sec. V.

Since we now have an explicit form for \tilde{V}_1 and \tilde{V}_2 in Eq. (36), we can reduce Eqs. (29), (30) and (31) to one dimensional integral equations. With the aid of the formulas given in the Appendix we obtain the following equations from Eqs. (29), (30) and (31):

$$\begin{aligned}
\tilde{A}(p_t^2) = & \frac{-1}{2W_p(E_0+m_D-W_p)} \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ \left[8\pi\kappa F_1(|p_t|, |q_t|) + \frac{32\pi\beta W_p}{3(Q_1^2-\mu^2)} [F_2(|p_t|, |q_t|, \mu) - F_2(|p_t|, |q_t|, Q_1)] \right] \tilde{A}(q_t^2) \right. \\
& - 8\pi\kappa F_1(|p_t|, |q_t|) \tilde{A}(p_t^2) \left. \right\} - \frac{1}{2W_p(E_0+m_D-W_p)} \int \frac{q_t^2 dq_t}{4\pi^2} \frac{8\pi\beta}{3(Q_1^2-\mu^2)} \left\{ q_t^2 [-F_2(|p_t|, |q_t|, \mu) + F_2(|p_t|, |q_t|, Q_1)] \right. \\
& - \frac{1}{p_t^2} [F_4(|p_t|, |q_t|, \mu) - F_4(|p_t|, |q_t|, Q_1)] \left. \right\} \tilde{C}(q_t^2) + \frac{1}{2W_p(E_0+m_D-W_p)} \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ 4\pi\kappa \left[q_t^2 F_1(|p_t|, |q_t|) \right. \right. \\
& - \frac{1}{p_t^2} F_5(|p_t|, |q_t|) \left. \right] + \frac{16\pi\beta W_p}{3(Q_1^2-\mu^2)} \left[q_t^2 [F_2(|p_t|, |q_t|, \mu) - F_2(|p_t|, |q_t|, Q_1)] + \frac{1}{2p_t^2} [F_4(|p_t|, |q_t|, \mu) \right. \\
& \left. \left. - F_4(|p_t|, |q_t|, Q_1)] \right] \right\} \tilde{D}(q_t^2), \tag{37}
\end{aligned}$$

$$\begin{aligned}
\tilde{C}(p_t^2) = & \frac{-1}{2m_D^2 W_p(E_0+m_D-W_p)} \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ \left[8\pi\kappa W_p F_1(|p_t|, |q_t|) + \frac{16\pi\beta}{3(Q_1^2-\mu^2)} [F_2(|p_t|, |q_t|, \mu) - F_2(|p_t|, |q_t|, Q_1)] [(E_0 \right. \right. \\
& \left. \left. + m_D) W_p + m_D^2] \right] \tilde{A}(q_t^2) - 8\pi\kappa W_p F_1(|p_t|, |q_t|) \tilde{A}(p_t^2) \right\} \\
& - \frac{1}{2m_D^2 W_p(E_0+m_D-W_p)} \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ \left[8\pi\kappa F_3(|p_t|, |q_t|) \frac{m_D^2 - (E_0+m_D)W_p}{p_t^2} + \frac{16\pi\beta}{3(Q_1^2-\mu^2)} [F_4(|p_t|, |q_t|, \mu) \right. \right. \\
& \left. \left. - F_4(|p_t|, |q_t|, Q_1)] \frac{W_p}{p_t^2} \right] \tilde{C}(q_t^2) - 8\pi\kappa F_1(|p_t|, |q_t|) [m_D^2 - (E_0+m_D)W_p] \tilde{C}(p_t^2) \right\} \\
& + \frac{1}{2m_D^2 W_p(E_0+m_D-W_p)} \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ \left[8\pi\kappa F_5(|p_t|, |q_t|) \frac{W_p}{p_t^2} - \frac{16\pi\beta}{3(Q_1^2-\mu^2)} [F_4(|p_t|, |q_t|, \mu) \right. \right. \\
& \left. \left. - F_4(|p_t|, |q_t|, Q_1)] \frac{(E_0+m_D)W_p + m_D^2}{p_t^2} \right] \tilde{D}(q_t^2) - 8\pi\kappa p_t^2 W_p F_1(|p_t|, |q_t|) \tilde{D}(p_t^2) \right\}, \tag{38}
\end{aligned}$$

$$\begin{aligned}
\tilde{D}(p_t^2) = & \frac{1}{2m_D^2 W_p(E_0+m_D-W_p)} \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ \left[8\pi\kappa F_1(|p_t|, |q_t|) + \frac{16\pi\beta W_p}{3(Q_1^2-\mu^2)} [F_2(|p_t|, |q_t|, \mu) - F_2(|p_t|, |q_t|, Q_1)] \right] \tilde{A}(q_t^2) \right. \\
& \left. - 8\pi\kappa F_1(|p_t|, |q_t|) \tilde{A}(p_t^2) \right\} + \frac{1}{2m_D^2 W_p(E_0+m_D-W_p)} \\
& \times \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ \left[-8\pi\kappa F_3(|p_t|, |q_t|) \frac{W_p}{p_t^2} + \frac{16\pi\beta}{3(Q_1^2-\mu^2)} [F_4(|p_t|, |q_t|, \mu) - F_4(|p_t|, |q_t|, Q_1)] \frac{2p_t^2 + 3m_D^2}{2p_t^4} \right. \right. \\
& \left. \left. + \frac{16\pi\beta}{3(Q_1^2-\mu^2)} [F_2(|p_t|, |q_t|, \mu) - F_2(|p_t|, |q_t|, Q_1)] \frac{q_t^2 m_D^2}{2p_t^2} \right] \tilde{C}(q_t^2) + 8\pi\kappa W_p F_1(|p_t|, |q_t|) \tilde{C}(p_t^2) \right\} \\
& - \frac{1}{2m_D^2 W_p(E_0+m_D-W_p)} \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ \left[8\pi\kappa F_5(|p_t|, |q_t|) \frac{2p_t^2 + 3m_D^2}{2p_t^4} - 8\pi\kappa F_1(|p_t|, |q_t|) \frac{q_t^2 m_D^2}{2p_t^2} \right. \right. \\
& \left. \left. - \frac{16\pi\beta}{3(Q_1^2-\mu^2)} [F_4(|p_t|, |q_t|, \mu) - F_4(|p_t|, |q_t|, Q_1)] \frac{W_p(p_t^2 + 3m_D^2)}{p_t^4} - \frac{16\pi\beta}{3(Q_1^2-\mu^2)} [F_2(|p_t|, |q_t|, \mu) \right. \right. \\
& \left. \left. - F_2(|p_t|, |q_t|, Q_1)] \frac{W_p m_D^2 q_t^2}{p_t^2} \right] \tilde{D}(q_t^2) - 8\pi\kappa (p_t^2 + m_D^2) F_1(|p_t|, |q_t|) \tilde{D}(p_t^2) \right\}, \tag{39}
\end{aligned}$$

where $|p_i| = \sqrt{p_i^2}$. The functions $F_i (i=1, \dots, 5)$, appearing in Eqs. (37), (38) and (39), are defined in the Appendix. From these three coupled integral equations we can solve numerically for $\tilde{A}(p_i^2)$, $\tilde{C}(p_i^2)$ and $\tilde{D}(p_i^2)$. This will be done in Sec. V. In the next section, we will first discuss the normalization of the BS wave function.

IV. NORMALIZATION FOR THE BS WAVE FUNCTION

It can be seen that the overall normalization of $\tilde{A}(p_i^2)$, $\tilde{C}(p_i^2)$ and $\tilde{D}(p_i^2)$ cannot be determined from Eqs. (37), (38) and (39). With the help of heavy quark symmetry, the normalization constant can be obtained from the fact that the Isgur-Wise function is normalized to 1 at the zero-recoil point. In the limit $m_Q \rightarrow \infty$ the weak transition matrix element induced by the current $\bar{c}\Gamma b$ for $\omega_b^{(*)} \rightarrow \omega_c^{(*)}$ has the following form from HQET:

$$\langle \omega_c^{(*)}(v') | \bar{c}\Gamma b | \omega_b^{(*)}(v) \rangle = \bar{B}_{m'}^\nu(v') \Gamma B_m^\mu(v) [\xi(\omega) g_{\mu\nu} + \zeta(\omega) v_\nu v'_\mu], \quad (40)$$

where $\omega = v \cdot v'$ is the velocity transfer, m, m' could be 1 or 2, and Γ is an arbitrary Dirac matrix. At the zero-recoil point, $v = v'$, only the $\xi(\omega) g_{\mu\nu}$ term contributes and we must have

$$\xi(\omega = 1) = 1. \quad (41)$$

On the other hand, the transition matrix element for $\omega_b^{(*)} \rightarrow \omega_c^{(*)}$ is related to the BS wave functions of $\omega_b^{(*)}$ and $\omega_c^{(*)}$ by the following equation:

$$\langle \omega_c^{(*)}(v') | \bar{c}\Gamma b | \omega_b^{(*)}(v) \rangle = \int \frac{d^4 p}{(2\pi)^4} \bar{\chi}_{P'm'}^\mu(p') \Gamma \chi_{Pm}^\nu(p) S_{D\mu\nu}^{-1}(p_2), \quad (42)$$

where $P (P')$ is the momentum of $\omega_b^{(*)}$ ($\omega_c^{(*)}$) and $\bar{\chi}_{P'm'}^\mu(p')$ is the wave function of the final state $\omega_c^{(*)}(v')$, which satisfies the constraint

$$\bar{\chi}_{P'm'}^\mu(p') \not{v}' = \bar{\chi}_{P'm'}^\mu(p'). \quad (43)$$

At the zero-recoil point, $p' = p$, since the light diquark sees no change in the heavy quark part, it does not change its relative momentum.

The scalar BS functions of the final state BS wave function obey the same BS equation as Eqs. (29), (30) and (31). Substituting Eq. (7) into Eq. (42) and using Eq. (40) we have

$$\xi(1) \bar{B}_{m'\mu}(v) \Gamma B_m^\mu(v) = \int \frac{d^4 p}{(2\pi)^4} \frac{i}{p_l + E_0 + m_D + i\epsilon} \bar{\chi}_{P'm'}^\mu(p) \Gamma \int \frac{d^4 q}{(2\pi)^4} G_{\mu\nu}(P, p, q) \chi_{Pm}^\nu(p). \quad (44)$$

Now we substitute the expression for the kernel, Eq. (18), and Eq. (17) into Eq. (44). Using the same technique as used for Eqs. (24), (25) and (26), we find that there is only the structure $\bar{B}_{m'\mu}(v) \Gamma B_{m\mu}(v)$ on the right hand side of Eq. (44). Substituting the explicit expressions for \tilde{V}_1 and \tilde{V}_2 into Eq. (36), and using the integration formulas in the Appendix, we arrive at the following expression for $\xi(1)$ after some tedious calculations:

$$\xi(1) = \int \frac{p_i^2 dp_i}{4\pi^2} \frac{2}{E_0 + m_D - W_p} \left[\tilde{A}(p_i^2) h_1(|p_i|) - \frac{1}{3} p_i^2 \tilde{C}(p_i^2) h_2(|p_i|) - \frac{1}{3} p_i^2 \tilde{D}(p_i^2) h_3(|p_i|) + \frac{1}{6m_D^2} p_i^2 h_2(|p_i|) h_4(|p_i|) \right], \quad (45)$$

where $h_i(|p_i|) (i=1,2,3,4)$ are given by the equations

$$\begin{aligned}
 h_1(|p_t|) = & \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ 8\pi\kappa F_1(|p_t|, |q_t|) \left[\tilde{A}(q_t^2) - \frac{1}{3} q_t^2 \tilde{D}(q_t^2) \right] + \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [F_2(|p_t|, |q_t|, \mu) - F_2(|p_t|, |q_t|, Q_1)] \right. \\
 & \left. \times \left[2W_p \tilde{A}(q_t^2) - \frac{1}{3} q_t^2 \tilde{C}(q_t^2) - \frac{2}{3} W_p q_t^2 \tilde{D}(q_t^2) \right] - 8\pi\kappa F_1(|p_t|, |q_t|) \left[\tilde{A}(p_t^2) - \frac{1}{3} p_t^2 \tilde{D}(p_t^2) \right] \right\}, \quad (46)
 \end{aligned}$$

$$\begin{aligned}
 h_2(|p_t|) = & \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [F_2(|p_t|, |q_t|, \mu) - F_2(|p_t|, |q_t|, Q_1)] \tilde{A}(q_t^2) + 8\pi\kappa F_3(|p_t|, |q_t|) \frac{1}{p_t^2} \tilde{C}(q_t^2) \right. \\
 & \left. + \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [F_4(|p_t|, |q_t|, \mu) - F_4(|p_t|, |q_t|, Q_1)] \frac{1}{p_t^2} \tilde{D}(q_t^2) - 8\pi\kappa F_1(|p_t|, |q_t|) \tilde{C}(p_t^2) \right\}, \quad (47)
 \end{aligned}$$

$$\begin{aligned}
 h_3(|p_t|) = & \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ 8\pi\kappa F_1(|p_t|, |q_t|) \tilde{A}(q_t^2) + \frac{32\pi\beta W_p}{3(Q_1^2 - \mu^2)} [F_2(|p_t|, |q_t|, \mu) - F_2(|p_t|, |q_t|, Q_1)] \tilde{A}(q_t^2) \right. \\
 & - 8\pi\kappa F_5(|p_t|, |q_t|) \frac{1}{p_t^2} \tilde{D}(q_t^2) + \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [F_4(|p_t|, |q_t|, \mu) - F_4(|p_t|, |q_t|, Q_1)] \frac{1}{p_t^2} [\tilde{C}(q_t^2) + 2W_p \tilde{D}(q_t^2)] \\
 & \left. - 8\pi\kappa F_1(|p_t|, |q_t|) [-\tilde{A}(p_t^2) + p_t^2 \tilde{D}(p_t^2)] \right\}, \quad (48)
 \end{aligned}$$

$$\begin{aligned}
 h_4(|p_t|) = & \int \frac{q_t^2 dq_t}{4\pi^2} \left\{ \frac{-16\pi\beta}{3(Q_1^2 - \mu^2)} [F_2(|p_t|, |q_t|, \mu) - F_2(|p_t|, |q_t|, Q_1)] \tilde{A}(q_t^2) + 8\pi\kappa F_3(|p_t|, |q_t|) \frac{1}{p_t^2} \tilde{C}(q_t^2) \right. \\
 & \left. - \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [F_4(|p_t|, |q_t|, \mu) - F_4(|p_t|, |q_t|, Q_1)] \frac{1}{p_t^2} \tilde{D}(q_t^2) - 8\pi\kappa F_1(|p_t|, |q_t|) \tilde{C}(p_t^2) \right\}. \quad (49)
 \end{aligned}$$

The BS scalar functions $\tilde{A}(p_t^2)$, $\tilde{C}(p_t^2)$ and $\tilde{D}(p_t^2)$ should be normalized such that they satisfy Eq. (45).

V. NUMERICAL SOLUTIONS FOR THE BS WAVE FUNCTION

In this section we solve the three coupled integral equations (37), (38) and (39) numerically. The method is to discretize the integration region into n pieces (with n sufficiently large). In this way, the integral equations become matrix equations and the BS scalar functions $\tilde{A}(p_t^2)$, $\tilde{C}(p_t^2)$ and $\tilde{D}(p_t^2)$ become n dimensional vectors. The matrix equations obtained in this way can be written in the following form:

$$\tilde{A} = Z_1 \tilde{A} + Z_2 \tilde{C} + Z_3 \tilde{D}, \quad (50)$$

$$R_1 \tilde{A} + R_2 \tilde{C} + R_3 \tilde{D} = 0, \quad (51)$$

$$T_1 \tilde{A} + T_2 \tilde{C} + T_3 \tilde{D} = 0, \quad (52)$$

where $Z_i, R_i, T_i (i=1,2,3)$ are $n \times n$ matrices and are given by Eqs. (37), (38) and (39).

Substituting Eqs. (51) and (52) into Eq. (50) we obtain the eigenvalue equation for \tilde{A} ,

$$H \tilde{A} = \tilde{A}, \quad (53)$$

where H is an $n \times n$ matrix:

$$\begin{aligned}
 H = & Z_1 + Z_2 (T_3^{-1} T_2 - R_3^{-1} R_2)^{-1} (R_3^{-1} R_1 - T_3^{-1} T_1) \\
 & + Z_3 (T_2^{-1} T_3 - R_2^{-1} R_3)^{-1} (R_2^{-1} R_1 - T_2^{-1} T_1). \quad (54)
 \end{aligned}$$

The eigenvalue equation (53) is solved by the so-called inverse iteration method [18]. In this way, we first construct the operator

$$K = \frac{1}{H - \lambda}, \quad (55)$$

where λ is some parameter which is chosen to be near to eigenvalue 1 in Eq. (53). In order to solve for the eigenvector \tilde{A} , we start with an arbitrary vector Y and operate K on Y sufficiently many times so that the eigenvector corresponding to the eigenvalue 1 dominates. In this way, the scalar function \tilde{A} is obtained.

TABLE I. Values of κ and $\alpha_s^{(\text{eff})}$ for $\Sigma_Q^{(*)}$ with three sets of m_D .

m_D (GeV)	0.90				
κ (GeV ³)	0.02	0.04	0.06	0.08	0.10
$\alpha_s^{(\text{eff})}$	0.5190	0.5593	0.5842	0.6061	0.6149
m_D (GeV)	0.95				
κ (GeV ³)	0.02	0.04	0.06	0.08	0.10
$\alpha_s^{(\text{eff})}$	0.5889	0.6123	0.6285	0.6406	0.6502
m_D (GeV)	1.0				
κ (GeV ³)	0.02	0.04	0.06	0.08	0.10
$\alpha_s^{(\text{eff})}$	0.6414	0.6560	0.6669	0.6757	0.6828

In our model we have several parameters, $\alpha_s^{(\text{eff})}$, κ , Q_1^2 , m_D and E_0 . The parameter Q_1^2 has been described in Sec. III, with $Q_1^2 = 3.2$ GeV² from the data of the electromagnetic form factor of the proton. It is noted that this value corresponds to the (qq') axial vector diquark ($q, q' = u$ or d), i.e., for $\Sigma_Q^{(*)}$. In the cases of $\Xi_Q^{(*)}$ or $\Omega_Q^{(*)}$ this value might be somewhat different because of $SU(3)$ flavor symmetry breaking. However, we do not have data to extract Q_1^2 for $\Xi_Q^{(*)}$ and $\Omega_Q^{(*)}$ at present. In this work, we simply use the same value for Q_1^2 based on the approximate $SU(3)$ flavor symmetry. On the other hand, the binding energy should satisfy the following relation:

$$m_{\omega_Q^{(*)}} = m_Q + m_D + E_0, \quad (56)$$

where we have omitted corrections of $O(1/m_Q)$, since we are working in the heavy quark limit. Note that $m_D + E_0$ is independent of the flavor of the heavy quark, because of the $SU(2)_f \times SU(2)_s$ symmetry. From the BS equation solutions in the meson case, it has been found that the values $m_b = 5.02$ GeV and $m_c = 1.58$ GeV give predictions which are in good agreement with experiments [15]. Hence in the baryon case we expect

$$m_D + E_0 = 0.88 \text{ GeV (for } \Sigma_Q^{(*)}), 1.07 \text{ GeV (for } \Xi_Q^{(*)}), \\ 1.12 \text{ GeV (for } \Omega_Q^{(*)}). \quad (57)$$

The parameter m_D cannot be determined, although there are suggestions from the analysis of valence structure functions that it should be around 0.9 GeV for non-strange diquarks [19]. Hence we let it vary within some reasonable range. When we solve the eigenvalue equation (53), the condition that the eigenvalue be 1 provides a relation between α_{seff} and κ . As discussed in Ref. [9] κ is related to κ' (κ' is the confinement parameter in the heavy meson case and is about 0.2 GeV² [14,15]), where $\kappa \sim \Lambda_{\text{QCD}} \kappa'$. Therefore, in our numerical calculations we let κ vary in the region between 0.02 GeV³ and 0.1 GeV³. The diquark mass, m_D , is chosen to vary from 0.9 GeV to 1 GeV for $\Sigma_Q^{(*)}$, from 1.1

TABLE II. Values of κ and $\alpha_s^{(\text{eff})}$ for $\Xi_Q^{(*)}$ with three sets of m_D .

m_D (GeV)	1.10				
κ (GeV ³)	0.02	0.04	0.06	0.08	0.10
$\alpha_s^{(\text{eff})}$	0.5047	0.5402	0.5643	0.5826	0.5974
m_D (GeV)	1.15				
κ (GeV ³)	0.02	0.04	0.06	0.08	0.10
$\alpha_s^{(\text{eff})}$	0.5785	0.5995	0.6155	0.6283	0.6391
m_D (GeV)	1.20				
κ (GeV ³)	0.02	0.04	0.06	0.08	0.10
$\alpha_s^{(\text{eff})}$	0.6341	0.6478	0.6588	0.6682	0.6763

GeV to 1.2 GeV for $\Xi_Q^{(*)}$, and from 1.15 GeV to 1.25 GeV for $\Omega_Q^{(*)}$. Then we obtain the parameter $\alpha_s^{(\text{eff})}$ for different values of m_D and κ . The numerical results are shown in Tables I, II and III for $\Sigma_Q^{(*)}$, $\Xi_Q^{(*)}$ and $\Omega_Q^{(*)}$, respectively.

With the parameters in Tables I, II and III we obtain the numerical solution for the BS scalar function \tilde{A} as the eigenvector of Eq. (53). Consequently, we get the numerical solutions for \tilde{C} and \tilde{D} from Eqs. (51) and (52). These solutions depend on the parameters m_D and κ . In Figs. 2, 3 and 4 we show the shapes of \tilde{A} , \tilde{C} and \tilde{D} for $\Sigma_Q^{(*)}$, $\Xi_Q^{(*)}$ and $\Omega_Q^{(*)}$ respectively. Figures 2(a), 3(a) and 4(a) show the dependence on κ for a typical m_D , while Figs. 2(b), 3(b) and 4(b) show the dependence on m_D for a typical κ . It can be seen from these figures that for different heavy baryons the shapes of the BS scalar functions are rather similar. This arises from the approximate $SU(3)$ flavor symmetry and is to be expected. All the scalar functions decrease to zero when $|p_i|$ is larger than about 1.5 GeV, because of the confinement interaction. Furthermore, since we are discussing the ground states $\omega_Q^{(*)}$, there are no nodes in the functions.

TABLE III. Values of κ and $\alpha_s^{(\text{eff})}$ for $\Omega_Q^{(*)}$ with three sets of m_D .

m_D (GeV)	1.15				
κ (GeV ³)	0.02	0.04	0.06	0.08	0.10
$\alpha_s^{(\text{eff})}$	0.4975	0.5325	0.5565	0.5748	0.5897
m_D (GeV)	1.20				
κ (GeV ³)	0.02	0.04	0.06	0.08	0.10
$\alpha_s^{(\text{eff})}$	0.5729	0.5935	0.6093	0.6222	0.6331
m_D (GeV)	1.25				
κ (GeV ³)	0.02	0.04	0.06	0.08	0.10
$\alpha_s^{(\text{eff})}$	0.6296	0.6430	0.6539	0.6633	0.6714




FIG. 2. The BS scalar wave functions for $\Sigma_Q^{(*)}$. The units are GeV^{-2} for $\bar{A}(p_i^2)$, GeV^{-3} for $\bar{C}(p_i^2)$ and GeV^{-4} for $\bar{D}(p_i^2)$. (a) For $m_D = 0.95$ GeV, we show the dependence on $|p_i|$ for two values of κ . In the upper plane, the upper [lower] solid line is for $\bar{A}(p_i^2)[\bar{C}(p_i^2)]$ with $\kappa = 0.02$ GeV^3 ; the upper [lower] dotted line is for $\bar{A}(p_i^2)[\bar{C}(p_i^2)]$ with $\kappa = 0.10$ GeV^3 . In the lower plane, the solid line is for $\bar{D}(p_i^2)$ with $\kappa = 0.02$ GeV^3 and the dotted line is for $\bar{D}(p_i^2)$ with $\kappa = 0.10$ GeV^3 . (b) For $\kappa = 0.06$ GeV^3 , we show the dependence on $|p_i|$ for two values of m_D . In the upper plane, the upper [lower] solid line is for $\bar{A}(p_i^2)[\bar{C}(p_i^2)]$ with $m_D = 0.90$ GeV; the upper [lower] dotted line is for $\bar{A}(p_i^2)[\bar{C}(p_i^2)]$ with $m_D = 1.00$ GeV. In the lower plane, the solid line is for $\bar{D}(p_i^2)$ with $m_D = 0.90$ GeV and the dotted line is for $\bar{D}(p_i^2)$ with $m_D = 1.00$ GeV.

VI. APPLICATION TO THE NONLEPTONIC DECAYS $\Omega_b \rightarrow \Omega_c^{(*)} P(V)$

In this section we will apply the numerical solutions of the BS equation to the nonleptonic decays $\Omega_b \rightarrow \Omega_c^{(*)}$ and a pseudoscalar or vector meson. In fact, $\Sigma_b^{(*)}$ and $\Xi_b^{(*)}$ decay strongly and their weak decays are hard to observe. However, Ω_b decays only weakly. We will first calculate the Isgur-Wise functions $\xi(\omega)$ and $\zeta(\omega)$ for $\Omega_b^{(*)} \rightarrow \Omega_c^{(*)}$ in Eq. (40) and then apply them to the nonleptonic weak decays of Ω_b .

A. Isgur-Wise functions for $\Omega_b^{(*)} \rightarrow \Omega_c^{(*)}$

The Isgur-Wise functions $\xi(\omega)$ and $\zeta(\omega)$ are related to the overlap integrals of the BS wave functions of the initial




FIG. 3. The BS scalar wave functions for $\Xi_Q^{(*)}$. The units are GeV^{-2} for $\bar{A}(p_i^2)$, GeV^{-3} for $\bar{C}(p_i^2)$ and GeV^{-4} for $\bar{D}(p_i^2)$. (a) For $m_D = 1.15$ GeV, we show the dependence on $|p_i|$ for two values of κ . In the upper plane, the upper [lower] solid line is for $\bar{A}(p_i^2)[\bar{C}(p_i^2)]$ with $\kappa = 0.02$ GeV^3 ; the upper [lower] dotted line is for $\bar{A}(p_i^2)[\bar{C}(p_i^2)]$ with $\kappa = 0.10$ GeV^3 . In the lower plane, the solid line is for $\bar{D}(p_i^2)$ with $\kappa = 0.02$ GeV^3 and the dotted line is for $\bar{D}(p_i^2)$ with $\kappa = 0.10$ GeV^3 . (b) For $\kappa = 0.06$ GeV^3 , we show the dependence on $|p_i|$ for two values of m_D . In the upper plane, the upper [lower] solid line is for $\bar{A}(p_i^2)[\bar{C}(p_i^2)]$ with $m_D = 1.10$ GeV; the upper [lower] dotted line is for $\bar{A}(p_i^2)[\bar{C}(p_i^2)]$ with $m_D = 1.20$ GeV. In the lower plane, the solid line is for $\bar{D}(p_i^2)$ with $m_D = 1.10$ GeV and the dotted line is for $\bar{D}(p_i^2)$ with $m_D = 1.20$ GeV.

(Ω_b) and final ($\Omega_c^{(*)}$) states. The concrete expression for them can be obtained by comparing the structure $\bar{B}_{m'}^\mu(v')\Gamma B_{m\mu}(v)$ and $v \cdot \bar{B}_{m'}(v')\Gamma v' \cdot B_m(v)$ on both sides of Eq. (40). Similarly to Eq. (44), we have the following equation after substituting Eq. (7) into Eq. (42) and using Eq. (40):

$$\begin{aligned} & \bar{B}_{m'}^\nu(v')\Gamma B_{m\mu}^\mu(v)[\xi(\omega)g_{\mu\nu} + \zeta(\omega)v_\nu v'_\mu] \\ &= \int \frac{d^4p}{(2\pi)^4} \frac{i}{p_l + E_0 + m_D + i\epsilon} \bar{\chi}_{Pm'}^\mu(p')\Gamma \\ & \times \int \frac{d^4q}{(2\pi)^4} G_{\mu\nu}(P, p, q)\chi_{Pm}^\nu(p). \end{aligned} \quad (58)$$




FIG. 4. The BS scalar wave functions for $\Omega_Q^{(*)}$. The units are GeV^{-2} for $\tilde{A}(p_t^2)$, GeV^{-3} for $\tilde{C}(p_t^2)$ and GeV^{-4} for $\tilde{D}(p_t^2)$. (a) For $m_D = 1.20$ GeV, we show the dependence on $|p_t|$ for two values of κ . In the upper plane, the upper [lower] solid line is for $\tilde{A}(p_t^2)[\tilde{C}(p_t^2)]$ with $\kappa = 0.02$ GeV^3 , the upper [lower] dotted line is for $\tilde{A}(p_t^2)[\tilde{C}(p_t^2)]$ with $\kappa = 0.10$ GeV^3 . In the lower plane, the solid line is for $\tilde{D}(p_t^2)$ with $\kappa = 0.02$ GeV^3 and the dotted line is for $\tilde{D}(p_t^2)$ with $\kappa = 0.10$ GeV^3 . (b) For $\kappa = 0.06$ GeV^3 , we show the dependence on $|p_t|$ for two values of m_D . In the upper plane, the upper [lower] solid line is for $\tilde{A}(p_t^2)[\tilde{C}(p_t^2)]$ with $m_D = 1.15$ GeV; the upper [lower] dotted line is for $\tilde{A}(p_t^2)[\tilde{C}(p_t^2)]$ with $m_D = 1.25$ GeV. In the lower plane, the solid line is for $\tilde{D}(p_t^2)$ with $m_D = 1.15$ GeV and the dotted line is for $\tilde{D}(p_t^2)$ with $m_D = 1.25$ GeV.

Substituting Eqs. (17) and (18) into Eq. (58) and using Eqs. (24), (25) and (26), we find that on the right hand side of Eq. (58) there are the following structures: $p'_t \cdot \bar{B}_{m'}(v') \Gamma v' \cdot B_m(v)$, $v \cdot \bar{B}_{m'}(v') \Gamma p_t \cdot B_m(v)$, $p'_t \cdot \bar{B}_{m'}(v') \Gamma p_t \cdot B_m(v)$, $p_t \cdot \bar{B}_{m'}(v') \Gamma p_t \cdot B_m(v)$, and $p'_t \cdot \bar{B}_{m'}(v') \Gamma p'_t \cdot B_m(v)$. However, all of them can be expressed in terms of $\bar{B}_{m'}^\mu(v') \Gamma B_{m\mu}(v)$ and $v \cdot \bar{B}_{m'}(v') \Gamma v' \cdot B_m(v)$, after integration over p , on the grounds of Lorentz invariance. Take $p'_t \cdot \bar{B}_{m'}(v') \Gamma p_t \cdot B_m(v)$ as an example. In general, the integral $\int [d^4 q / (2\pi)^4] p_t'^\mu p_t^\nu f$, where f is some Lorentz scalar function, can be expressed in terms of $g^{\mu\nu}$, $v^\mu v^\nu$, $v'^\mu v'^\nu$, $v'^\mu v^\nu$ and $v^\mu v'^\nu$. However, only

the $g^{\mu\nu}$ and $v^\mu v'^\nu$ terms contribute when contracted with $\bar{B}_{m'}^\mu(v') \Gamma B_{m\nu}(v)$, leading to the structures $\bar{B}_{m'}^\mu(v') \Gamma B_{m\mu}(v)$ and $v \cdot \bar{B}_{m'}(v') \Gamma v' \cdot B_m(v)$, respectively. The coefficients of these two terms can be obtained directly. In this way, we have the following replacement rule:

$$\begin{aligned}
& p'_t \cdot \bar{B}_{m'}(v') \Gamma v' \cdot B_m(v) \\
& \rightarrow \frac{1}{1-\omega^2} v \cdot p'_t v \cdot \bar{B}_{m'}(v') \Gamma v' \cdot B_m(v), \\
& v \cdot \bar{B}_{m'}(v') \Gamma p_t \cdot B_m(v) \\
& \rightarrow \frac{1}{1-\omega^2} v' \cdot p_t v \cdot \bar{B}_{m'}(v') \Gamma v' \cdot B_m(v), \\
& p'_t \cdot \bar{B}_{m'}(v') \Gamma p_t \cdot B_m(v) \\
& \rightarrow \left[\frac{1}{2} p_t \cdot p'_t - \frac{\omega}{2(\omega^2-1)} v \cdot p'_t v' \cdot p_t \right] \bar{B}_{m'}^\mu(v') \Gamma B_{m\mu}(v) \\
& + \left[-\frac{\omega}{2(\omega^2-1)} p_t \cdot p'_t + \frac{\omega^2+2}{2(\omega^2-1)^2} v \cdot p'_t v' \cdot p_t \right] \\
& \times v \cdot \bar{B}_{m'}(v') \Gamma v' \cdot B_m(v), \\
& p_t \cdot \bar{B}_{m'}(v') \Gamma p_t \cdot B_m(v) \\
& \rightarrow \left[-\frac{1}{2} p_t^2 + \frac{1}{2(\omega^2-1)} (v' \cdot p_t)^2 \right] \bar{B}_{m'}^\mu(v') \Gamma B_{m\mu}(v) \\
& + \left[\frac{\omega}{2(\omega^2-1)} p_t^2 - \frac{3\omega}{2(\omega^2-1)^2} (v' \cdot p_t)^2 \right] \\
& \times v \cdot \bar{B}_{m'}(v') \Gamma v' \cdot B_m(v), \\
& p'_t \cdot \bar{B}_{m'}(v') \Gamma p'_t \cdot B_m(v) \\
& \rightarrow \left[-\frac{1}{2} p_t'^2 + \frac{1}{2(\omega^2-1)} (v \cdot p_t')^2 \right] \bar{B}_{m'}^\mu(v') \Gamma B_{m\mu}(v) \\
& + \left[\frac{\omega}{2(\omega^2-1)} p_t'^2 - \frac{3\omega}{2(\omega^2-1)^2} (v \cdot p_t')^2 \right] \\
& \times v \cdot \bar{B}_{m'}(v') \Gamma v' \cdot B_m(v). \tag{59}
\end{aligned}$$

Since in the weak transition the diquark acts as a spectator, its momentum in the initial and final baryons should be the same, $p_2 = p_2'$. Then we can show that

$$p' = p + m_D(v' - v), \tag{60}$$

where we have omitted the $O(1/m_Q)$ corrections. From Eq. (60) we have the following relations straightforwardly:

$$p'_t = p_t \omega - p_t \sqrt{\omega^2 - 1} \cos \theta,$$

$$\begin{aligned}
 p_t'^2 &= p_t^2 + p_t^2(\omega^2 - 1)\cos^2 \theta + p_t^2(\omega^2 - 1) - 2p_t p_t \omega \sqrt{\omega^2 - 1} \cos \theta, \\
 v' \cdot p_t &= -p_t \sqrt{\omega^2 - 1} \cos \theta, \\
 v \cdot p_t' &= (1 - \omega^2)p_t + \omega p_t \sqrt{\omega^2 - 1} \cos \theta, \\
 p_t \cdot p_t' &= -p_t^2 - p_t^2(\omega^2 - 1)\cos^2 \theta + \omega \sqrt{\omega^2 - 1} p_t p_t \cos \theta,
 \end{aligned} \tag{61}$$

where θ is the angle between p_t and v_t' .

With the aid of the relations between $A(p), C(p), D(p)$ and $\tilde{A}(p_t^2), \tilde{C}(p_t^2), \tilde{D}(p_t^2)$ [Eqs. (32)–(34)] and using Eqs. (59), (61) and the integration formulas in the Appendix we have the explicit expressions for $\xi(\omega)$ and $\zeta(\omega)$ after integrating the p_t component by selecting the proper contour,

$$\begin{aligned}
 \xi(\omega) &= \int \frac{p_t^2 dp_t}{4\pi^2} \int_0^\pi \sin \theta d\theta \frac{-1}{2W_p(E_0 + m_D - W_p)(E_0 + m_D - \omega W_p - p_t \sqrt{\omega^2 - 1} \cos \theta)} \\
 &\times \left\{ -2W_p(E_0 + m_D - W_p)F_A(p_t^2, \cos \theta)\tilde{A}(p_t^2) - \frac{3}{4}(1 - \cos^2 \theta)[h_1(|p_t|) - h_3(|p_t|)]F_A(p_t^2, \cos \theta) \right. \\
 &- \frac{1}{2}\omega p_t^2(1 - \cos^2 \theta)F_C(p_t^2, \cos \theta)h_2(|p_t|) + \frac{3}{4}p_t \sqrt{\omega^2 - 1}(1 - \cos^2 \theta)\cos \theta[h_1(|p_t|) - h_3(|p_t|)]F_C(p_t^2, \cos \theta) + W_p(1 \\
 &- \cos^2 \theta)(E_0 + m_D - W_p)p_t^2\tilde{A}(p_t^2)F_D(p_t^2, \cos \theta) - \frac{1}{2}p_t^2(1 - \cos^2 \theta)[(\omega^2 - 1)W_p \\
 &+ \omega p_t \sqrt{\omega^2 - 1} \cos \theta]h_2(|p_t|)F_D(p_t^2, \cos \theta) + \frac{3}{4}(1 - \cos^2 \theta)[p_t^2 + p_t^2(\omega^2 - 1)\cos^2 \theta + \omega \sqrt{\omega^2 - 1}W_p p_t]F_D(p_t^2, \cos \theta) \\
 &\left. \times [h_1(|p_t|) - h_3(|p_t|)] \right\},
 \end{aligned} \tag{62}$$

and

$$\begin{aligned}
 \zeta(\omega) &= \int \frac{p_t^2 dp_t}{4\pi^2} \int_0^\pi \sin \theta d\theta \frac{-1}{2W_p(E_0 + m_D - W_p)(E_0 + m_D - \omega W_p - p_t \sqrt{\omega^2 - 1} \cos \theta)} \left\{ \frac{1}{\sqrt{\omega^2 - 1}} p_t \cos \theta F_A(p_t^2, \cos \theta) h_2(|p_t|) \right. \\
 &+ \frac{3\omega}{4(\omega^2 - 1)} (1 - 3\cos^2 \theta)[h_1(|p_t|) - h_3(|p_t|)]F_A(p_t^2, \cos \theta) + \left[W_p + \frac{\omega}{\sqrt{\omega^2 - 1}} p_t \cos \theta \right] \\
 &\times F_C(p_t^2, \cos \theta) 2W_p(E_0 + m_D - W_p)\tilde{A}(p_t^2) + \frac{\omega}{2(\omega^2 - 1)} [\omega p_t^2 - 3\omega \cos^2 \theta p_t^2 - 2W_p p_t \sqrt{\omega^2 - 1} \cos \theta] \\
 &\times F_C(p_t^2, \cos \theta) h_2(|p_t|) - \frac{3}{4\sqrt{\omega^2 - 1}} [\omega p_t - 3\omega \cos^2 \theta p_t - 2W_p \sqrt{\omega^2 - 1} \cos \theta] \cos \theta [h_1(|p_t|) - h_3(|p_t|)] \\
 &\times F_C(p_t^2, \cos \theta) + \left[-\omega W_p^2 - \frac{2\omega^2}{\sqrt{\omega^2 - 1}} W_p p_t \cos \theta - \frac{\omega(2\omega^2 + 1)}{2(\omega^2 - 1)} p_t^2 \cos^2 \theta + \frac{\omega}{2(\omega^2 - 1)} p_t^2 \right] \\
 &\times [-2W_p(E_0 + m_D - W_p)]F_D(p_t^2, \cos \theta)\tilde{A}(p_t^2) + \frac{1}{2\sqrt{\omega^2 - 1}} [\omega p_t^2 - 3\omega \cos^2 \theta p_t^2 - 2W_p p_t \sqrt{\omega^2 - 1} \cos \theta] \\
 &\times (\sqrt{\omega^2 - 1}W_p + \omega p_t \cos \theta)F_D(p_t^2, \cos \theta)h_2(|p_t|) - \frac{3}{4(\omega^2 - 1)} [\omega p_t - 3\omega \cos^2 \theta p_t - 2W_p \sqrt{\omega^2 - 1} \cos \theta] \\
 &\times [p_t + p_t(\omega^2 - 1) \\
 &\times \cos^2 \theta + \omega \sqrt{\omega^2 - 1}W_p \cos \theta]F_D(p_t^2, \cos \theta)[h_1(|p_t|) - h_3(|p_t|)] \left. \right\},
 \end{aligned} \tag{63}$$

where $h_i(|p_t|)$ ($i=1,2,3,4$) are given in Eqs. (46)–(49) and $F_A(p_t^2, \cos \theta), F_C(p_t^2, \cos \theta)$ and $F_D(p_t^2, \cos \theta)$ have the following expressions:

$$\begin{aligned}
F_A(p_i^2, \cos \theta) = & \int \frac{q_i^2 dq_i}{4\pi^2} \left\{ 8\pi\kappa F_1(|p_i'|, |q_i|) \left[\tilde{A}(q_i^2) - \frac{1}{2} q_i^2 \tilde{D}(q_i^2) - \tilde{A}(p_i'^2) \right] + \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [F_2(|p_i'|, |q_i|, \mu) \right. \\
& - F_2(|p_i'|, |q_i|, Q_1)] \left[2(\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta) \left(\tilde{A}(q_i^2) - \frac{1}{2} q_i^2 \tilde{D}(q_i^2) \right) - \frac{1}{2} q_i^2 \tilde{C}(q_i^2) \right] \\
& + \frac{1}{2p_i'^2} \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [-F_4(|p_i'|, |q_i|, \mu) + F_4(|p_i'|, |q_i|, Q_1)] [\tilde{C}(q_i^2) + 2(\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta) \tilde{D}(q_i^2)] \\
& \left. + \frac{1}{2p_i'^2} 8\pi\kappa F_5(|p_i'|, |q_i|) \tilde{D}(q_i^2) \right\}, \tag{64}
\end{aligned}$$

$$\begin{aligned}
F_C(p_i^2, \cos \theta) = & \frac{1}{m_D^2} \int \frac{q_i^2 dq_i}{4\pi^2} \left\{ 8\pi\kappa F_1(|p_i'|, |q_i|) \{ (\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta) [\tilde{A}(q_i^2) - \tilde{A}(p_i'^2) + p_i'^2 \tilde{D}(p_i'^2)] \right. \\
& + [(\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta)^2 - m_D^2] \tilde{C}(p_i'^2) \} + \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [F_2(|p_i'|, |q_i|, \mu) - F_2(|p_i'|, |q_i|, Q_1)] \\
& \times [(\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta)^2 + m_D^2] \tilde{A}(q_i^2) - \frac{1}{p_i'^2} 8\pi\kappa F_3(|p_i'|, |q_i|) [(\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta)^2 - m_D^2] \\
& \times \tilde{C}(q_i^2) - \frac{1}{p_i'^2} \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [-F_4(|p_i'|, |q_i|, \mu) + F_4(|p_i'|, |q_i|, Q_1)] \{ (\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta) \tilde{C}(q_i^2) \\
& \left. + [(\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta)^2 + m_D^2] \tilde{D}(q_i^2) \} - \frac{1}{p_i'^2} 8\pi\kappa F_5(|p_i'|, |q_i|) (\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta) \tilde{D}(q_i^2) \right\}, \tag{65}
\end{aligned}$$

$$\begin{aligned}
F_D(p_i^2, \cos \theta) = & -\frac{1}{m_D^2} \int \frac{q_i^2 dq_i}{4\pi^2} \left\{ 8\pi\kappa F_1(|p_i'|, |q_i|) \left[\tilde{A}(q_i^2) + \frac{m_D^2}{2p_i'^2} q_i^2 \tilde{D}(q_i^2) - \tilde{A}(p_i'^2) \right. \right. \\
& \left. + (\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta) \tilde{C}(p_i'^2) + (p_i'^2 + m_D^2) \tilde{D}(p_i'^2) \right] + \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [F_2(|p_i'|, |q_i|, \mu) \\
& - F_2(|p_i'|, |q_i|, Q_1)] \left[(\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta) \left(\tilde{A}(q_i^2) + \frac{m_D^2}{p_i'^2} q_i^2 \tilde{D}(q_i^2) \right) + \frac{m_D^2}{2p_i'^2} q_i^2 \tilde{C}(q_i^2) \right] \\
& - \frac{1}{p_i'^2} 8\pi\kappa F_3(|p_i'|, |q_i|) (\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta) \tilde{C}(q_i^2) - \frac{1}{p_i'^2} \frac{16\pi\beta}{3(Q_1^2 - \mu^2)} [-F_4(|p_i'|, |q_i|, \mu) \\
& + F_4(|p_i'|, |q_i|, Q_1)] \left[\frac{3m_D^2 + 2p_i'^2}{2p_i'^2} [\tilde{C}(q_i^2) + 2(\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta) \tilde{D}(q_i^2)] \right. \\
& \left. - (\omega W_p + p_i \sqrt{\omega^2 - 1} \cos \theta) \tilde{D}(q_i^2) \right] - \frac{3m_D^2 + 2p_i'^2}{2p_i'^2} 8\pi\kappa F_5(|p_i'|, |q_i|) \tilde{D}(q_i^2) \left. \right\}. \tag{66}
\end{aligned}$$

In Sec. V we obtained the numerical results for $\tilde{A}(p_i^2)$, $\tilde{C}(p_i^2)$ and $\tilde{D}(p_i^2)$. Substituting these results into Eqs. (62) and (63) we have the numerical solutions for $\xi(\omega)$ and $\zeta(\omega)$ depending on the parameters in our model. For $\Omega_b^{(*)}$

$\rightarrow \Omega_c^{(*)}$, we show the Isgur-Wise functions with typical value $m_D = 1.20$ GeV in Fig. 5(a) ($\kappa = 0.02$ GeV³) and Fig. 5(b) ($\kappa = 0.10$ GeV³) respectively. The dependence of the Isgur-Wise functions on m_D is shown in Fig. 5(c) (m_D




FIG. 5. (a)–(d) Numerical solutions for $\xi(\omega)$ and $\zeta(\omega)$ for $\Omega_b^{(*)} \rightarrow \Omega_c^{(*)}$. The upper solid line is $\xi(\omega)$ and the lower solid line is $\zeta(\omega)$. The dotted line is $-\xi(\omega)/(\omega+1)$. The parameters are $m_D=1.20$ GeV and $\kappa=0.02$ GeV³ in (a), $m_D=1.20$ GeV and $\kappa=0.10$ GeV³ in (b), $\kappa=0.06$ GeV³ and $m_D=1.15$ GeV in (c), $\kappa=0.06$ GeV³ and $m_D=1.25$ GeV in (d).

$=1.15$ GeV) and Fig. 5(d) ($m_D=1.25$ GeV) for $\kappa=0.06$ GeV³. It can be seen from these plots that $\xi(\omega)$ and $\zeta(\omega)$ have opposite signs and $\xi(\omega)$ changes more rapidly than $\zeta(\omega)$ as ω increases.

It is interesting to study the relation between $\xi(\omega)$ and $\zeta(\omega)$. Based on the picture that in the large N_c limit heavy baryons are viewed as the bound states of chiral solitons and heavy mesons [20], Chow has shown that $\xi(\omega)$ and $\zeta(\omega)$ obey the following relation [21]:

$$\xi(\omega) = -(1+\omega)\zeta(\omega). \quad (67)$$

The deviation from this relation is caused by $1/N_c$ corrections. From Figs. 5(a)–(d) we can see that, in the range of the parameters in our model, Eq. (67) is generally satisfied. For some sets of parameters this relation holds well.

B. Nonleptonic decays $\Omega_b \rightarrow \Omega_c^{(*)} P(V)$

In this section we will discuss the Cabibbo-allowed two body nonleptonic decays of $\Omega_b \rightarrow \Omega_c^{(*)} P(V)$ (P and V stand for pseudoscalar and vector mesons respectively). The Hamiltonian describing such decays reads

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* (a_1 O_1 + a_2 O_2), \quad (68)$$

with $O_1 = (\bar{D}U)(\bar{c}b)$ and $O_2 = (\bar{c}U)(\bar{D}b)$, where U and D are the fields for light quarks involved in the decay, and $(\bar{q}_1 q_2) = \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2$ is understood. The parameters a_1 and a_2 are treated as free parameters since they involve hadronization effects. Since Ω_b decays are energetic, the factorization assumption is applied so that one of the currents in the Hamiltonian (68) is factorized out and generates a meson [22,23]. Thus the decay amplitude of the two body nonleptonic decay becomes the product of two matrix elements; one is related to the decay constant of the factorized meson (P or V) and the other is the weak transition matrix element between Ω_b and $\Omega_c^{(*)}$,

$$M^{\text{fac}}(\Omega_b \rightarrow \Omega_c^{(*)} P(V)) = \frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 \langle P(V) | A_\mu(V_\mu) | 0 \rangle \times \langle \Omega_c^{(*)}(v') | J^\mu | \Omega_b(v) \rangle. \quad (69)$$

Here J_μ is the weak current $(\bar{c}b)$ and $\langle 0 | A_\mu(V_\mu) | P(V) \rangle$ are related to the decay constants of the pseudoscalar or vector mesons by

$$\begin{aligned} \langle 0 | A_\mu | P \rangle &= i f_P q_\mu, \\ \langle 0 | V_\mu | V \rangle &= f_V m_V \epsilon_\mu, \end{aligned} \quad (70)$$

where q_μ is the momentum of the emitted meson (from the W boson), ϵ_μ is the polarization vector of the vector meson and the normalization for the decay constants is chosen so that $f_\pi = 132$ MeV. It is noted that in the two body nonleptonic weak decays $\Omega_b \rightarrow \Omega_c^{(*)} P(V)$ there is no contribution from the a_2 term, since such a term corresponds to the transition of Ω_b to a light baryon instead of $\Omega_c^{(*)}$. On the other hand, the general form for the amplitudes of $\Omega_b \rightarrow \Omega_c^{(*)} P(V)$ is

$$\begin{aligned} M(\Omega_b \rightarrow \Omega_c P) &= i\bar{u}_f(v') (A + B\gamma_5) u_i(v), \\ M(\Omega_b \rightarrow \Omega_c V) &= \bar{u}_f(v') \epsilon^{*\mu} [A_1 \gamma_\mu \gamma_5 + A_2 p_{f\mu} \gamma_5 \\ &\quad + B_1 \gamma_\mu + B_2 p_{f\mu}] u_i(v), \\ M(\Omega_b \rightarrow \Omega_c^* P) &= i q_\mu \bar{u}_f^\mu(v') (C + D\gamma_5) u_i(v), \\ M(\Omega_b \rightarrow \Omega_c^* V) &= \bar{u}_f^\nu(v') \epsilon^{*\mu} [g_{\nu\mu} (C_1 + D_1 \gamma_5) \\ &\quad + p_{i\nu} \gamma_\mu (C_2 + D_2 \gamma_5) + p_{i\nu} p_{f\mu} (C_3 \\ &\quad + D_3 \gamma_5)] u_i(v), \end{aligned} \quad (71)$$

where u_i is the Dirac spinor of Ω_b , $u_f^{(\mu)}$ is the Dirac (Rarita-Schwinger) spinor of $\Omega_c^{(*)}$, and $p_{i(f)}$ is the momentum of Ω_b ($\Omega_c^{(*)}$).

From Eqs. (69)–(71) and using Eq. (40) we find

$$\begin{aligned} A &= \frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_P \frac{1}{3} (m_i - m_f) [(\omega + 2)\xi(\omega) \\ &\quad + (\omega^2 - 1)\zeta(\omega)], \\ B &= \frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_P \frac{1}{3} (m_i + m_f) [(3\omega - 2)\xi(\omega) \\ &\quad + 3(\omega^2 - 1)\zeta(\omega)], \\ A_1 = B_1 &= \frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_V m_V \frac{1}{3} [\omega\xi(\omega) \\ &\quad + (\omega^2 - 1)\zeta(\omega)], \\ A_2 &= \frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_V m_V \frac{2}{3} \left(\frac{1}{m_i} - \frac{1}{m_f} \right) \\ &\quad \times [\xi(\omega) + (\omega + 1)\zeta(\omega)], \\ B_2 &= -\frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_V m_V \frac{2}{3} \left(\frac{1}{m_i} + \frac{1}{m_f} \right) \\ &\quad \times [\xi(\omega) + (\omega - 1)\zeta(\omega)], \\ C &= -\frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_P \frac{1}{\sqrt{3}} \left(1 + \frac{m_f}{m_i} \right) \\ &\quad \times [\xi(\omega) + (\omega - 1)\zeta(\omega)], \end{aligned}$$

$$\begin{aligned} D &= -\frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_P \frac{1}{\sqrt{3}} \left[\left(1 - \frac{m_f}{m_i} \right) \xi(\omega) \right. \\ &\quad \left. + \left(\omega - 1 - (\omega + 3) \frac{m_f}{m_i} \right) \zeta(\omega) \right], \\ C_1 = D_1 &= \frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_V m_V \frac{2}{\sqrt{3}} \xi(\omega), \\ C_2 &= -\frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_V m_V \frac{1}{\sqrt{3}} \frac{1}{m_i} \\ &\quad \times [\xi(\omega) + (\omega + 1)\zeta(\omega)], \\ D_2 &= \frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_V m_V \frac{1}{\sqrt{3}} \frac{1}{m_i} \\ &\quad \times [\xi(\omega) + (\omega - 1)\zeta(\omega)], \\ C_3 = D_3 &= \frac{G_F}{\sqrt{2}} V_{cb} V_{UD}^* a_1 f_V m_V \frac{2}{\sqrt{3}} \frac{1}{m_i m_f} \zeta(\omega), \end{aligned} \quad (72)$$

where m_i (m_f) is the mass of Ω_b ($\Omega_c^{(*)}$).

With Eqs. (71) and (72) we can calculate the decay widths and polarization parameters for $\Omega_b \rightarrow \Omega_c^{(*)} P(V)$. The kinematic formulas which have been derived using both partial wave and helicity methods can be found in Refs. [24,25]. These two methods are equivalent. For instance, in the helicity method [25], the decay width is expressed in terms of the helicity amplitudes,

$$\Gamma = \frac{p_c}{16\pi m_i^2} \sum_{\lambda_i, \lambda_f} |h_{\lambda_f, \lambda_{P(V)}; \lambda_i}|^2, \quad (73)$$

where p_c is the c.m. momentum of the decay products and the helicity amplitudes are defined as

$$\begin{aligned} h_{\lambda_f, \lambda_{P(V)}; \lambda_i} &= \langle \Omega_c^{(*)}(\lambda_f), P(V)(\lambda_{P(V)}) | H_{\text{eff}} | \Omega_b(\lambda_i) \rangle \\ &= (\lambda_f - \lambda_{P(V)} = \lambda_i). \end{aligned} \quad (74)$$

The ‘‘up-down’’ asymmetry is given by

$$\alpha = \frac{\sum_{\lambda_f} (|h_{\lambda_f, \lambda_{P(V)}; 1/2}|^2 - |h_{\lambda_f, \lambda_{P(V)}; -1/2}|^2)}{\sum_{\lambda_i, \lambda_f} |h_{\lambda_f, \lambda_{P(V)}; \lambda_i}|^2}. \quad (75)$$

The relations between the helicity amplitudes and the amplitudes given in Eq. (71), which we will not list here, can be found in [25,26]. Then from Eqs. (72)–(75), we obtain the numerical results for the decay widths and asymmetry parameters. In Table IV we list the results for $m_D = 1.20$ GeV. The numbers without (with) brackets correspond to $\kappa = 0.02$ GeV³ ($\kappa = 0.10$ GeV³). The results for κ

TABLE IV. Predictions for decay widths and asymmetry parameters for $\Omega_b^- \rightarrow \Omega_c^{(*)} P(V)$ for $m_D = 1.20$ GeV.

Process	Γ (10^{10} s^{-1})	α
$\Omega_b^- \rightarrow \Omega_c^0 \pi^-$	$0.052a_1^2$ ($0.154a_1^2$)	-0.67 (-0.70)
$\Omega_b^- \rightarrow \Omega_c^0 D_s^-$	$0.261a_1^2$ ($0.592a_1^2$)	-0.56 (-0.58)
$\Omega_b^- \rightarrow \Omega_c^0 \rho^-$	$0.073a_1^2$ ($0.207a_1^2$)	-0.68 (-0.71)
$\Omega_b^- \rightarrow \Omega_c^0 D_s^{*-}$	$0.115a_1^2$ ($0.245a_1^2$)	-0.73 (-0.74)
$\Omega_b^- \rightarrow \Omega_c^{*0} \pi^-$	$0.046a_1^2$ ($0.133a_1^2$)	-0.61 (-0.58)
$\Omega_b^- \rightarrow \Omega_c^{*0} D_s^-$	$0.165a_1^2$ ($0.370a_1^2$)	-0.54 (-0.52)
$\Omega_b^- \rightarrow \Omega_c^{*0} \rho^-$	$0.134a_1^2$ ($0.354a_1^2$)	0.59 (0.59)
$\Omega_b^- \rightarrow \Omega_c^{*0} D_s^{*-}$	$0.462a_1^2$ ($0.960a_1^2$)	0.31 (0.31)

$= 0.06 \text{ GeV}^3$ in the range $m_D = 1.15 \text{ GeV}$ (without brackets) and $m_D = 1.25 \text{ GeV}$ (with brackets) are shown in Table V. In the calculations we have taken $m_{\Omega_b} = 6.14 \text{ GeV}$ and the following decay constants:

$$f_\pi = 132 \text{ MeV}, \quad f_{D_s} = 241 \text{ MeV} [27], \quad f_\rho = 216 \text{ MeV},$$

$$f_{D_s} = f_{D_s^*}.$$

It can be seen from Tables IV and V that the predictions for the decay widths show a strong dependence on the parameters κ and m_D in our model. The experimental data in the future will be used to fix these parameters and test our model. However, the dependence of the up-down asymmetries on these parameters is slight.

The decay widths and asymmetry parameters have also been calculated in the nonrelativistic quark model approach [26], where the form factors are calculated at the zero-recoil point and then extrapolated to other ω values under the assumption of a dipole behavior. By comparing the predictions in the BS and quark models we find that the decay widths in

 TABLE V. Predictions for decay widths and asymmetry parameters for $\Omega_b^- \rightarrow \Omega_c^{(*)} P(V)$ for $\kappa = 0.06 \text{ GeV}^3$.

Process	Γ (10^{10} s^{-1})	α
$\Omega_b^- \rightarrow \Omega_c^0 \pi^-$	$0.075a_1^2$ ($0.145a_1^2$)	-0.64 (-0.72)
$\Omega_b^- \rightarrow \Omega_c^0 D_s^-$	$0.358a_1^2$ ($0.562a_1^2$)	-0.54 (-0.59)
$\Omega_b^- \rightarrow \Omega_c^0 \rho^-$	$0.102a_1^2$ ($0.150a_1^2$)	-0.65 (-0.70)
$\Omega_b^- \rightarrow \Omega_c^0 D_s^{*-}$	$0.149a_1^2$ ($0.232a_1^2$)	-0.71 (-0.75)
$\Omega_b^- \rightarrow \Omega_c^{*0} \pi^-$	$0.067a_1^2$ ($0.123a_1^2$)	-0.64 (-0.56)
$\Omega_b^- \rightarrow \Omega_c^{*0} D_s^-$	$0.227a_1^2$ ($0.345a_1^2$)	-0.56 (-0.50)
$\Omega_b^- \rightarrow \Omega_c^{*0} \rho^-$	$0.200a_1^2$ ($0.314a_1^2$)	0.59 (0.59)
$\Omega_b^- \rightarrow \Omega_c^{*0} D_s^{*-}$	$0.616a_1^2$ ($0.888a_1^2$)	0.31 (0.31)

our model are smaller than those in the quark model. For the asymmetry parameters, the difference is even larger. Except for the processes $\Omega_b^- \rightarrow \Omega_c^{*0} \pi^-$ and $\Omega_b^- \rightarrow \Omega_c^{*0} D_s^-$ in Tables IV and V, even the signs of α in these two models are opposite.

VII. SUMMARY AND DISCUSSION

Since in the heavy quark limit the light degrees of freedom in a heavy baryon have good spin and isospin quantum numbers and since the internal structure is blind to the flavor and spin direction of the heavy quark, we assume that a heavy baryon, $\omega_Q^{(*)}$, is composed of a heavy quark and a light axial vector diquark. Based on this picture, we establish the BS equation for the heavy baryon $\omega_Q^{(*)}$. We discuss the form of the BS wave function and find that in the heavy quark limit there are three BS scalar functions to describe the dynamics inside a heavy baryon $\omega_Q^{(*)}$. This is consistent with our physical picture. In order to solve the BS equation, we assume a kernel containing a scalar confinement term and a one-gluon-exchange term, as in the Λ_Q case. In the heavy quark limit, the heavy quark is almost on massshell inside a heavy baryon and it is appropriate to apply the covariant instantaneous approximation in the kernel. Then we derive explicitly three coupled integral equations for the three BS scalar functions \tilde{A} , \tilde{C} and \tilde{D} . These equations are solved numerically and we give the model predictions for these functions. The results appear reasonable. It is shown that the shapes of these functions are similar for $\Sigma_Q^{(*)}$, $\Xi_Q^{(*)}$ and $\Omega_Q^{(*)}$, with differences arising from $SU(3)$ flavor symmetry breaking effects.

Although the BS equation is formally the exact equation to describe the bound state, there is much difficulty in applying it to the real physical state. The most difficult point is that we must take a phenomenologically inspired form for the kernel. Furthermore, we have used the quark and diquark propagators with their free form, which leads to some uncertainties. In our approach, there are several parameters such as κ , m_D and α_s^{eff} , subject to the condition that the observed masses of $\omega_Q^{(*)}$ be reproduced. In our numerical solutions we let these parameters vary in some reasonable range. Another parameter is Q_1^2 , which arises from the internal structure of diquark. Its value for the (qq') diquark ($q, q' = u$ or d) is extracted from the data of the electromagnetic form factor of the proton. When there is a strange quark in the diquark, we do not have a means to determine its exact value at present. In the future, the experimental data for $\omega_Q^{(*)}$ should help to fix the parameters in our model.

As phenomenological applications, this model has been used to calculate the the Isgur-Wise functions $\xi(\omega)$ and $\zeta(\omega)$ for $\Omega_b^{(*)} \rightarrow \Omega_c^{(*)}$, and consequently, has provided theoretical predictions for the Cabibbo-allowed two body non-leptonic decay rates and up-down asymmetries for the physical processes $\Omega_b \rightarrow \Omega_c^{(*)} P(V)$. It has been shown that the relation between $\xi(\omega)$ and $\zeta(\omega)$ in our model is generally consistent with that in the soliton model in the large N_c limit. We have also compared our model predictions with those in the nonrelativistic quark model. Our model yields decay

widths which are much smaller and for the asymmetry parameters the difference is even bigger. All these predictions will be tested in future experiments.

ACKNOWLEDGMENTS

This work was supported in part by the Australian Research Council and the National Science Foundation of China.

APPENDIX: INTEGRATION FORMULAS

In this appendix we give the formulas which are used to reduce the three dimensional integration to the one dimensional integration. In the following formulas $\phi(q_i^2)$ is some arbitrary function of q_i^2 . The relevant results needed are

$$\begin{aligned} I_1 &= \int \frac{d^3 q_i}{(2\pi)^3} \frac{\phi(q_i^2)}{[(p_i - q_i)^2 + \mu^2]^2} \\ &= \int \frac{q_i^2 dq_i}{4\pi^2} \phi(q_i^2) F_1(|p_i|, |q_i|), \end{aligned} \quad (\text{A1})$$

with

$$F_1(|p_i|, |q_i|) = \frac{2}{(p_i^2 + q_i^2 + \mu^2)^2 - 4p_i^2 q_i^2}; \quad (\text{A2})$$

$$\begin{aligned} I_2 &= \int \frac{d^3 q_i}{(2\pi)^3} \frac{\phi(q_i^2)}{(p_i - q_i)^2 + \delta^2} \\ &= \int \frac{q_i^2 dq_i}{4\pi^2} \phi(q_i^2) F_2(|p_i|, |q_i|, \delta), \end{aligned} \quad (\text{A3})$$

with

$$F_2(|p_i|, |q_i|, \delta) = \frac{1}{2|p_i||q_i|} \ln \frac{(|p_i| + |q_i|)^2 + \delta^2}{(|p_i| - |q_i|)^2 + \delta^2}; \quad (\text{A4})$$

$$\begin{aligned} I_3 &= \int \frac{d^3 q_i}{(2\pi)^3} \frac{p_i \cdot q_i \phi(q_i^2)}{[(p_i - q_i)^2 + \mu^2]^2} \\ &= \int \frac{q_i^2 dq_i}{4\pi^2} \phi(q_i^2) F_3(|p_i|, |q_i|), \end{aligned} \quad (\text{A5})$$

with

$$\begin{aligned} F_3(|p_i|, |q_i|) &= \frac{1}{4|p_i||q_i|} \left[\ln \frac{(|p_i| - |q_i|)^2 + \mu^2}{(|p_i| + |q_i|)^2 + \mu^2} \right. \\ &\quad \left. + \frac{4|p_i||q_i|(p_i^2 + q_i^2 + \mu^2)}{(p_i^2 + q_i^2 + \mu^2)^2 - 4p_i^2 q_i^2} \right]; \end{aligned} \quad (\text{A6})$$

$$\begin{aligned} I_4 &= \int \frac{d^3 q_i}{(2\pi)^3} \frac{(p_i \cdot q_i)^2 \phi(q_i^2)}{(p_i - q_i)^2 + \delta^2} \\ &= - \int \frac{q_i^2 dq_i}{4\pi^2} \phi(q_i^2) F_4(|p_i|, |q_i|, \delta), \end{aligned} \quad (\text{A7})$$

with

$$\begin{aligned} F_4(|p_i|, |q_i|, \delta) &= \frac{p_i^2 + q_i^2 + \delta^2}{2} \left[1 + \frac{p_i^2 + q_i^2 + \delta^2}{4|p_i||q_i|} \right. \\ &\quad \left. \times \ln \frac{(|p_i| - |q_i|)^2 + \delta^2}{(|p_i| + |q_i|)^2 + \delta^2} \right]; \end{aligned} \quad (\text{A8})$$

$$\begin{aligned} I_5 &= \int \frac{d^3 q_i}{(2\pi)^3} \frac{(p_i \cdot q_i)^2 \phi(q_i^2)}{[(p_i - q_i)^2 + \mu^2]^2} \\ &= \int \frac{q_i^2 dq_i}{4\pi^2} \phi(q_i^2) F_5(|p_i|, |q_i|), \end{aligned} \quad (\text{A9})$$

with

$$\begin{aligned} F_5(|p_i|, |q_i|) &= \frac{1}{2} \left[1 + \frac{p_i^2 + q_i^2 + \mu^2}{2|p_i||q_i|} \ln \frac{(|p_i| - |q_i|)^2 + \mu^2}{(|p_i| + |q_i|)^2 + \mu^2} \right. \\ &\quad \left. + \frac{(p_i^2 + q_i^2 + \mu^2)^2}{(p_i^2 + q_i^2 + \mu^2)^2 - 4p_i^2 q_i^2} \right]. \end{aligned} \quad (\text{A10})$$

- [1] N. Isgur and M.B. Wise, Phys. Lett. B **232**, 113 (1989); **237**, 527 (1990); H. Georgi, *ibid.* **264**, 447 (1991); see also M. Neubert, Phys. Rep. **245**, 259 (1994) for the review.
[2] OPAL Collaboration, R. Akers *et al.*, Z. Phys. C **69**, 195 (1996); Phys. Lett. B **353**, 402 (1995); OPAL Collaboration, K. Ackerstaff *et al.*, *ibid.* **426**, 161 (1998).
[3] UA1 Collaboration, C. Albarjar *et al.*, Phys. Lett. B **273**, 540 (1991).

- [4] CDF Collaboration, F. Abe *et al.*, Phys. Rev. D **47**, 2639 (1993).
[5] S. E. Tzmaris, in *High Energy Physics*, Proceedings of the 27th International Conference on High Energy Physics, Scotland, 1994, edited by P. J. Bussey and I. G. Knowles (IOP, Bristol, 1995); P. Abreu *et al.*, Phys. Lett. B **374**, 351 (1996).
[6] CDF Collaboration, F. Abe *et al.*, Phys. Rev. D **55**, 1142 (1997).

- [7] Particle Data Group, C. Caso *et al.*, Eur. Phys. J. C **3**, 1 (1998).
- [8] T. Mannel, W. Roberts and Z. Ryzak, Nucl. Phys. **B355**, 38 (1991).
- [9] X.-H. Guo and T. Muta, Phys. Rev. D **54**, 4629 (1996); Mod. Phys. Lett. A **11**, 1523 (1996).
- [10] A. Falk, Nucl. Phys. **B378**, 79 (1992).
- [11] T. Mannel, W. Roberts and Z. Ryzak, Phys. Lett. B **271**, 421 (1991); W. Roberts, Nucl. Phys. **B389**, 549 (1993).
- [12] Y.-B. Dai, X.-H. Guo and C.-S. Huang, Nucl. Phys. **B412**, 277 (1994).
- [13] D. Lurie, *Particles and Fields* (Wiley, New York, 1968); C. Itzykson and J.B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).
- [14] E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane and T.-M. Yan, Phys. Rev. D **17**, 3090 (1978).
- [15] H.-Y. Jin, C.-S. Huang and Y.-B. Dai, Z. Phys. C **56**, 707 (1992); Y.-B. Dai, C.-S. Huang and H.-Y. Jin, *ibid.* **60**, 527 (1993).
- [16] M. Anselmino, P. Kroll and B. Pire, Z. Phys. C **36**, 89 (1987); P. Kroll, B. Quadder and W. Schweiger, Nucl. Phys. **B316**, 373 (1988).
- [17] G.P. Lepage and S.J. Brodsky, Phys. Rev. D **22**, 2157 (1980); S. J. Brodsky, G. P. Lepage, T. Huang and P.B. MacKenzie, in *Particles and Fields 2*, edited by A. Z. Capri and A. N. Kamal (Plenum, New York, 1983), p. 83.
- [18] Y. Rae Kwon and F. Tabakin, Phys. Rev. C **18**, 932 (1978); C. Savkli and F. Tabakin, Nucl. Phys. **A628**, 645 (1998).
- [19] F. Close and A.W. Thomas, Phys. Lett. B **212**, 227 (1988).
- [20] E. Jenkins, A.V. Manohar and M.B. Wise, Nucl. Phys. **B396**, 27 (1993); **B396**, 38 (1993); Z. Guralnik, M. Luke and A.V. Manohar, *ibid.* **B390**, 474 (1993).
- [21] C.-K. Chow, Phys. Rev. D **51**, 1224 (1995); **54**, 873 (1996).
- [22] J.D. Bjorken, Nucl. Phys. B (Proc. Suppl.) **11**, 325 (1989).
- [23] M.J. Dugan and B. Grinstein, Phys. Lett. B **255**, 583 (1991).
- [24] S. Pakvasa, S.F. Tuan and S.P. Rosen, Phys. Rev. D **42**, 3746 (1990).
- [25] J.G. Körner and M. Krämer, Z. Phys. C **55**, 659 (1992).
- [26] H.-Y. Cheng, Phys. Rev. D **56**, 2799 (1997).
- [27] C.T. Sachrajda, in *Beauty 96*, Rome, Italy, 1996 [Nucl. Instrum. Methods Phys. Res. A **384**, 26 (1996)].