THE PETROLOGY, GEOCHEMISTRY, AND FLUID HISTORY OF CALC-SILICATE ROCKS AND ASSOCIATED PRIMARY Cu-Co MINERALIZATION IN THE ETHIUDNA MINES AREA, OLARY PROVINCE, SOUTH AUSTRALIA.

Benjamin Grguric, BSc.

Thesis submitted as partial fulfilment of the Honours Degree of Bachelor of Science

Department of Geology and Geophysics, University of Adelaide

November, 1992

NATIONAL GRID REFERENCE: PLUMBAGO SHEET 6833 (1:63,360)

COPY2

TABLE OF CONTENTS

CHAPTER 1	
INTRODUCTION	1
1.1. INTRODUCTION	1
1.2. HISTORY AND PREVIOUS INVESTIGATIONS	1
1.3. AIMS AND METHODS OF STUDY	2
CHAPTER 2	
GEOLOGICAL BACKGROUND	3
2.1. REGIONAL GEOLOGY AND STRATIGRAPHY OF THE	
WILLYAMA SUPERGROUP	
2.2. STRATIGRAPHY OF THE ETHIUDNA SUB-BLOCK	5
2.2.1. Biotite Microclinolite	5
2.2.2. Metapelites	6
2.2.3. Ethiudna Adamellite	6
2.3. STRUCTURAL GEOLOGY OF THE ETHIUDNA SUB-BLOCK	7
2.4. METAMORPHISM OF THE ETHIUDNA SUB-BLOCK	8
CHAPTER 3	
PETROLOGY AND FLUID HISTORY OF THE BIMBA SUITE	10
3.1. INTRODUCTION	10
3.2. PETROGRAPHY OF THE BIMBA SUITE	
3.2.1. The Wollastonite-Diopside Subunit	10
3.2.2. The Calc-Silicate Quartzite	
3.2.3. The Footwall Quartzite	12
3.3. PETROGENESIS OF THE BIMBA SUITE	12
3.3.1 Description of textures	12
3.3.2 Interpretation of textures	
implications for fluid infiltration	13
3.4. AQUEOUS ELECTROLYTE COMPOSITIONS OF	
METASOMATIC FLUIDS	16
3.4.1. Fluid inclusion analysis	
3.4.2. Scapolite analysis	18
3.4.3. Origin of high metamorphic fluid salinities	21
3.5 CARBON-OXYGEN STABLE ISOTOPE GEOCHEMISTRY OF	
THE BIMBA SUITE	2 1
CHAPTER 4	
SULPHIDE PETROLOGY AND GENESIS	2 5
4.1. THE NATURE AND DISTRIBUTION OF PRIMARY	
SULPHIDE MINERALIZATION	2 5
4.1.1. Ethiudna mines	
4.1.2. Ethiudna East prospect	
4.1.3. Other sulphide occurrences	
4.2. ORE TEXTURES AND MINERALOGY	27

4.3. DISTRIBUTION AND STRUCTURAL CONTROL OF
MINERALIZATION
GENETIC IMPLICATIONS3
4.4. GENESIS OF STRATA-BOUND Cu-Co MINERALIZATION3 1
4.5. EVIDENCE FOR METASOMATIC REMOBILIZATION OF
MINERALIZATION3 3
ACKNOWLEDGEMENTS
REFERENCES
APPENDIX 1: WHOLE ROCK ANALYSES
APPENDIX 2: CARBON-OXYGEN ISOTOPE ANALYSES
APPENDIX 3: MICROPROBE ANALYSES-SCAPOLITE
APPENDIX 4: MICROPROBE ANALYSES-SULPHIDES
APPENDIX 5: FLUID INCLUSION ANALYSES
APPENDIX 6: SULPHUR ISOTOPE ANALYSES
APPENDIX 7: UNIVARIANT REACTIONS AND SLOPES OF UNIVARIANT LINES

LIST OF FIGURES

Figure 1	Location of the study area.
Figure 2	Schematic stratigraphy of the Willyama Supergroup.
	metasediments in the study area, and regional correlations.
Figures 3a and	13b
_	Discriminant diagrams for altered volcanics and
	volcanoclastics.
Figure 4	XCO2-T diagram showing univariant curves for the reaction
V.	calcite + quartz => wollastonite + CO ₂ .
Figure 5	μK2O-μCO2 diagram for the system KCASH.
Figure 6	Histograms of data from microthermometric measurements
	on fluid inclusions.
Figure 7	Compositions of Ethiudna scapolites.
Figure 8	Correlations between scapolite composition and fluid
	composition.
Figure 9	Carbon-oxygen isotope profiles.
Figure 10	Sulphide species proportions: stratigraphic variation.
Figure 11	T-X diagram for phase relations in the system Fe-S.
Figure 12	Ternary plot of Ethiudna pyrite compositions.
Figure 13	Histogram of sulphur isotope analyses.
Figure 14	Drill hole assay profiles.
	LIST OF TABLES
Table 1	Modal compositions of Ethiudna microclinolites.
Table 2	Wollastonite Stage assemblages.
Table 3	
	Carbon-oxygen isotope analyses. Mineralogy of sulphide mineralization.
Table 4	Geochemical characteristics of Ethiudna mineralization.
Table 5	Geochemical characteristics of Eunuchia mineralization.
	LIST OF PLATES
Plates 1-6	Photographs and photomicrographs displaying textural
	relationships in calc-silicates.
Plates 7-8	Photomicrographs of fluid inclusions.
Plates 9-17	Photomicrographs of ore textures.

ABSTRACT

Textures in calc-silicate rocks of the Lower Proterozoic Bimba Suite in the Ethiudna mines area, Olary Province, South Australia, record multiple fluid infiltrational episodes associated with both prograde, amphibolite-facies, and retrograde, greenschist-facies metamorphism. Pervasive infiltration of H₂O-rich fluids (XCO₂<0.18) during prograde metamorphism, was associated with the large-scale development of wollastonite-rich lithologies within the Bimba Suite. Stable oxygen isotope evidence suggests these fluids were equilibrated with a siliciclastic sequence(s) prior to infiltrating the Bimba Suite.

A second, retrograde metamorphic (temperatures ~300 °C) infiltrational episode was associated with the development of a grossular-quartz assemblage. Fluid flow was channelized during this episode, the H₂O-rich fluids being focussed along structurally-induced zones of high permeability. Large variations in fluid/rock ratios are recorded in the calc-silicates as large variations in the progress of the grossular-quartz producing reaction.

The formation of cross-cutting quartz, calcite, and rare laumontite veins postdates both these infiltrational episodes. Fluid inclusion evidence suggests these veins were deposited by fluids at temperatures of less than 250°C.

Fluid inclusion data and scapolite compositional data indicate fluids associated with all three episodes were highly saline, consistent with conclusions made by earlier workers that the Bimba Suite represents a meta-evaporite sequence.

The microcline-rich rocks of the Quartzofeldspathic Suite, which underlies the Bimba Suite, host stratiform and disseminated Cu-Co sulphide mineralization. Geochemical and petrological evidence suggests this mineralization is of the genetic type known as a red-bed-associated Cu deposit. The distribution, geochemistry, and textural characteristics of sulphides in the Bimba Suite, suggest this mineralization represents a metasomatic remobilization of pre-metamorphic Quartzofeldspathic Suite mineralization. The sulphide-silicate-carbonate textural relationship in the Bimba Suite suggests this remobilization took place during the high temperature, wollastonite-producing infiltrational episode, and that precipitation of ore metals was effected by a pH rise.