

**The genetic improvement of wheat and barley for reproductive  
frost tolerance**

**By**

**Jason Reinheimer**

**Bachelor of Agricultural Science, University of Adelaide**

**A thesis submitted for the degree of Doctor of Philosophy in the Faculty of  
Sciences at the University of Adelaide**

**School of Agriculture, Food and Wine  
Waite Campus**

**2010**

## **Bibliography**

Allard, W. A. (1999) Principles of plant breeding, John Wiley and Sons.

Andrews, C. J., Pomeroy, M. K. & Roche, L. A. D. L. (1973) Cold-acclimation of winter-wheat during early seedling growth. *Plant Physiology*, 51, 26-26.

Baga, M., Chodaparambil, S. V., Limin, A. E., Pecar, M., Fowler, D. B. & Chibbar, R. N. (2007) Identification of quantitative trait loci and associated candidate genes for low-temperature tolerance in cold-hardy winter wheat. *Functional and Integrative Genomics*, 7, 53-68.

Beck, E. H., Fettig, S., Knake, C., Hartig, K. & Bhattarai, T. (2007) Specific and unspecific responses of plants to cold and drought stress. *Journal of Biosciences*, 32, 501-510.

Borner, A., Buck-Sorlin, G. H., Hayes, P. M., Malyshev, S. & Korzun, V. (2002) Molecular mapping of major genes and quantitative trait loci determining flowering time in response to photoperiod in barley. *Plant Breeding*, 121, 129-132.

Brougham, M. (2006) The relationship between vernalisation sensitivity, photoperiod sensitivity, basic vegetative period and grain yield in Australian wheat germplasm. Honours Thesis. Discipline of Plant and Pest Science. Adelaide, University of Adelaide.

Campbell, S. A. & Close, T. J. (1997) Dehydrins: genes, proteins, and associations with phenotypic traits. *New Phytologist*, 137, 61-74.

- Cattivelli, L., Baldi, P., Crosatti, C., Di Fonzo, N., Faccioli, P., Grossi, M., Mastrangelo, A. M., Pecchioni, N. & Stanca, A. M. (2002a) Chromosome regions and stress-related sequences involved in resistance to abiotic stress in Triticeae. *Plant Molecular Biology*, 48, 649-665.
- Cattivelli, L., Crosatti, C., Mare, C., Grossi, M., Mastrangelo, A. M., Mazzucotelli, C. G., Galiba, G. & Stanca, A. M. (2002b) Expression of cold-regulated (*cor*) genes in barley: Molecular basis and environmental interaction. IN Li, P. H. & Palva, E. T. (Eds.) *Plant Cold Hardiness: Gene Regulation and Genetic Engineering*. New York, Kluwer Academic/Plenum Publishers.
- Chen, A., Brule-Babel, A., Baumann, U. & Collins, N. (2009a) Structure-function analysis of the barley genome: the gene-rich region of chromosome 2HL. *Functional and Integrative Genomics*, 9, 67-79.
- Chen, A., Reinheimer, J., Brule-Babel, A., Baumann, U., Pallotta, M., Fincher, G. & Collins, N. (2009b) Genes and traits associated with chromosome 2H and 5H regions controlling sensitivity of reproductive tissues to frost in barley. *Theoretical and Applied Genetics*, 118, 1465-1476.
- Chen, Y. H., Carver, B. F., Wang, S. W., Zhang, F. Q. & Yan, L. L. (2009c) Genetic loci associated with stem elongation and winter dormancy release in wheat. *Theoretical and Applied Genetics*, 118, 881-889.
- Choi, D. W., Rodriguez, E. M. & Close, T. J. (2002) Barley *Cbf3* gene identification, expression pattern, and map location. *Plant Physiology*, 129, 1781-1787.

- Choi, D. W., Zhu, B. & Close, T. J. (1999) The barley (*Hordeum vulgare* L.) dehydrin multigene family: sequences, allele types, chromosome assignments, and expression characteristics of 11 *Dhn* genes of cv Dicktoo. *Theoretical and Applied Genetics*, 98, 1234-1247.
- Collins, H. M., Panozzo, J. F., Logue, S. J., Jefferies, S. P. & Barr, A. R. (2003) Mapping and validation of chromosome regions associated with high malt extract in barley (*Hordeum vulgare* L.). *Australian Journal of Agricultural Research*, 54, 1223-1240.
- Coventry, S. J., Barr, A. R., Eglinton, J. K. & McDonald, G. K. (2003) The determinants and genome locations influencing grain weight and size in barley (*Hordeum vulgare* L.). *Australian Journal of Agricultural Research*, 54, 1103-1115.
- Coventry, S. J. & Eglinton, J. K. (2005) Collaborative barley breeding for low rainfall environments: Harvest Report 1999-2004. University of Adelaide.
- Eglinton, J. K. (2003) Novel alleles from wild barley for breeding malting barley (*Hordeum vulgare* L.). Ph.D Thesis. School of Agriculture, Food and Wine : Plant and Pest Science Adelaide, University of Adelaide.
- Fletcher, R. J. (1988) Relationship between freezing injury suffered during stem elongation stage of development and vernalization response in a winter wheat cultivar (*Triticum aestivum* L. cv. Cheyenne). *Australian Journal of Agricultural Research*, 39, 129-140.
- Foote, T. N., Griffiths, S., Allouis, S. & Moore, G. (2004) Construction and analysis of a BAC library in the grass *Brachypodium sylvaticum*: its use as a tool to bridge the gap

between rice and wheat in elucidating gene content. *Functional and Integrative Genomics*, 4, 26-33.

Fowler, D. B., Chauvin, L.P., Limin, A.E., Sarhan, F. (1996) The regulatory role of vernalization in the expression of low-temperature-induced genes in wheat and rye. *Theoretical and Applied Genetics*, 93, 554-559.

Fowler, D. B. & Limin, A. E. (2004) Interactions among factors regulating phenological development and acclimation rate determine low-temperature tolerance in wheat. *Annals of Botany*, 94, 717-724.

Fowler, D. B., Limin, A.E., Gusta, L.V. (1983) Breeding for winter hardiness in wheat. IN Fowler, D. B., Gusta, L.V., Sinklard, A.E., Hobin, B.A. (Ed.) *New frontiers of winter wheat production*. Saskatoon, University of Saskatchewan.

Fox, G. P., Panozzo, J. F., Li, C. D., Lance, R. C. M., Inkerman, P. A. & Henry, R. J. (2003) Molecular basis of barley quality. *Australian Journal of Agricultural Research*, 54, 1081-1101.

Francia, E., Barabaschi, D., Tondelli, A., Laido, G., Rizza, F., Stanca, A. M., Busconi, M., Fogher, C., Stockinger, E. J. & Pecchioni, N. (2007) Fine mapping of a *HvCBF* gene cluster at the frost resistance locus *Fr-H2* in barley. *Theoretical and Applied Genetics*, 115, 1083-1091.

Francia, E., Rizza, F., Cattivelli, L., Stanca, A. M., Galiba, G., Toth, B., Hayes, P. M., Skinner, J. S. & Pecchioni, N. (2004) Two loci on chromosome 5H determine low-

temperature tolerance in a 'Nure' (winter) x 'Tremois' (spring) barley map. *Theoretical and Applied Genetics*, 108, 670-680.

Fredericks, T., Christophers, J. & Borrell, A. (2005) Investigation of post head-emergence frost resistance of barley in the northern cropping region of Australia. 12th Australian Barley Technical Symposium, 1-3.

Frisch, M. & Melchinger, A. E. (2005) Selection theory for marker-assisted backcrossing. *Genetics*, 170, 909-917.

Fu, D. L., Szucs, P., Yan, L. L., Helguera, M., Skinner, J. S., Von Zitzewitz, J., Hayes, P. M. & Dubcovsky, J. (2005) Large deletions within the first intron in *VRN-1* are associated with spring growth habit in barley and wheat. *Molecular Genetics and Genomics*, 273, 54-65.

Fuller, M. P., Fuller, A. M., Kaniouras, S., Christophers, J. & Fredericks, T. (2007) The freezing characteristics of wheat at ear emergence. *European Journal of Agronomy*, 26, 435-441.

Galiba, G., Kerepesi, I., Vagujfalvi, A., Kocsy, G., Cattivelli, L., Dubcovsky, J., Snape, J.W. (2001) Mapping of genes involved in glutathione, carbohydrate and *COR14b* cold induced protein accumulation during cold hardening in wheat. *Euphytica*, 119, 173-177.

Galiba, G., Quarrie, S.A., Sutka, J. And Morgounov, A. (1995) RFLP mapping of the vernalization (*Vrn1*) and frost resistance (*Fr1*) genes on chromosome 5A of wheat. *Theoretical and Applied Genetics*, 90, 1174-1179.

- Hayes, P. M., Blake, T., Chen, T. H. H., Tragoonrung, S., Chen, F., Pan, A. & Liu, B. (1993) Quantitative trait loci on barley (*Hordeum vulgare* L.) chromosome 7 associated with components of winterhardiness. *Genome*, 36, 66-71.
- Hearnden, P. R., Eckermann, P. J., McMichael, G. L., Hayden, M. J., Eglinton, J. K. & Chalmers, K. J. (2007) A genetic map of 1,000 SSR and DArT markers in a wide barley cross. *Theoretical and Applied Genetics*, 115, 383-391.
- Hoekstra, S., Vanzijderveld, M. H., Louwse, J. D., Heidekamp, F. & Vandermark, F. (1992) Anther and microspore culture of *Hordeum vulgare* L. cv Igri. *Plant Science*, 86, 89-96.
- Hospital, F. (2001) Size of donor chromosome segments around introgressed loci and reduction of linkage drag in marker-assisted backcross programs. *Genetics*, 158, 1363-1379.
- Huner, N. P. A., Oquist, G., Hurry, V. M., Krol, M., Falk, S. & Griffith, M. (1993) Photosynthesis, photoinhibition and low temperature acclimation in cold tolerant plants. *Photosynthesis Research*, 37, 19-39.
- Jefferies, S. P. (2000) Marker assisted backcrossing for gene introgression in barley. Ph. D. Department of Plant Science. Adelaide, University of Adelaide.
- Kalberer, S. R., Wisniewski, M. & Arora, R. (2006) Deacclimation and reacclimation of cold-hardy plants: Current understanding and emerging concepts. *Plant Science*, 171, 3-16.

- Karakousis, A., Barr, A. R., Kretschmer, J. M., Manning, S., Logue, S. J., Roumeliotis, S., Collins, H. M., Chalmers, K. J., Li, C. D., Lance, R. C. M. & Langridge, P. (2003a) Mapping and QTL analysis of the barley population Galleon x Haruna Nijo. *Australian Journal of Agricultural Research*, 54, 1131-1135.
- Karakousis, A., Gustafson, J. P., Chalmers, K. J., Barr, A. R. & Langridge, P. (2003b) A consensus map of barley integrating SSR, RFLP, and AFLP markers. *Australian Journal of Agricultural Research*, 54, 1173-1185.
- Karsai, I., Meszaros, K., Hayes, P. M. & Bedo, Z. (1997) Effects of loci on chromosomes 2 (2H) and 7 (5H) on developmental patterns in barley (*Hordeum vulgare* L.) under different photoperiod regimes. *Theoretical and Applied Genetics*, 94, 612-618.
- Koemel, J. E., Guenzi, A. C., Anderson, J. A. & Smith, E. L. (2004) Cold hardiness of wheat near-isogenic lines differing in vernalization alleles. *Theoretical and Applied Genetics*, 109, 839-846.
- Kosambi, D. D. (1944) The estimation of map distance from recombination values. *Annals of Eugenics*, 12, 172-175.
- Kota, R., Rudd, S., Facius, A., Kolesov, G., Thiel, T., Zhang, H., Stein, N., Mayer, K. & Graner, A. (2003) Snipping polymorphisms from large EST collections in barley (*Hordeum vulgare* L.). *Molecular Genetics and Genomics*, 270, 24-33.
- Koti, K., Karsai, I., Szucs, P., Horvath, C., Meszaros, K., Kiss, G. B., Bedo, Z. & Hayes, P. M. (2006) Validation of the two-gene epistatic model for vernalization response in a winter x spring barley cross. *Euphytica*, 152, 17-24.



- Laurie, D. A., Pratchett, N., Benzant, J. H. & Snape, J. W. (1995a) RFLP mapping of five major genes and eight quantitative trait loci controlling flowering time in a winter x spring barley (*Hordeum vulgare* L.) cross. *Genome*, 38, 575-585.
- Laurie, D. A., Pratchett, N., Bezant, J. & Snape, J. W. (1995b) RFLP mapping of 13 loci controlling flowering time in a winter x spring barley (*Hordeum vulgare* L.) cross. *Genome*, 38, 575-585.
- Limin, A. E. & Fowler, D. B. (2006) Low-temperature tolerance and genetic potential in wheat (*Triticum aestivum* L.): response to photoperiod, vernalization, and plant development. *Planta*, 224, 360-366.
- Lindelaursen, I., Heslopharrison, J. S., Shepherd, K. W. & Taketa, S. (1997) The barley genome and its relationship with the wheat genomes. A survey with an internationally agreed recommendation for barley chromosome nomenclature. *Hereditas*, 126, 1-16.
- Loukoianov, A., Yan, L. L., Blechl, A., Sanchez, A. & Dubcovsky, J. (2005) Regulation of *VRN-1* vernalization genes in normal and transgenic polyploid wheat. *Plant Physiology*, 138, 2364-2373.
- Mahfoozi, S., Limin, A. E. & Fowler, D. B. (2001) Influence of vernalization and photoperiod responses on cold hardiness in winter cereals. *Crop Science*, 41, 1006-1011.
- Marcellos, H. (1988) Comments on frost injury in wheat. *Frost Injury of Wheat - A Workshop of the Wheat Research Council*, 100-107.

- Marcellos, H. & Single, W. V. (1976) Ice nucleation on wheat. *Agricultural Meteorology*, 16, 125-129.
- McDonald, G., Eglinton, J. & Barr, A. (2009) Assessment of the agronomic value of QTL on chromosomes 2H and 4H linked to tolerance to boron toxicity in barley (*Hordeum vulgare* L.). *Plant and Soil*, 275-290.
- McDonald, G. K., Sutton, B. G. & Ellison, F. W. (1983) The effect of time of sowing on the grain yield of irrigated wheat in the Namoi Valley, New South Wales. *Australian Journal of Agricultural Research*, 34, 229-240.
- Pallotta, M. A., Asayama, S., Reinheimer, J. M., Davies, P. A., Barr, A. R., Jefferies, S. P., Chalmers, K. J., Lewis, J., Collins, H. M., Roumeliotis, S., Logue, S. J., Coventry, S. J., Lance, R. C. M., Karakousis, A., Lim, P., Verbyla, A. P. & Eckermann, P. J. (2003) Mapping and QTL analysis of the barley population Amagi Nijo x WI2585. *Australian Journal of Agricultural Research*, 54, 1141-1144.
- Pan, A., Hayes, P. M., Chen, F., Blake, T., Wright, S., Karsai, I. & Bedö, Z. (1994) Genetic analysis of the components of winterhardiness in barley (*Hordeum vulgare* L.). *Theoretical and Applied Genetics*, 89, 900-910.
- Park, R. F., Poulsen, D., Barr, A. R., Cakir, M., Moody, D. B., Raman, H. & Read, B. J. (2003) Mapping genes for resistance to *Puccinia hordei* in barley. *Australian Journal of Agricultural Research*, 54, 1323-1333.
- Pidal, B., Yan, L. L., Fu, D. L., Zhang, F. Q., Tranquilli, G. & Dubcovsky, J. (2009) The CArG-box located upstream from the transcriptional start of wheat vernalization

gene *VRN1* Is not necessary for the vernalization response. *Journal of Heredity*, 100, 355-364.

Prasil, I. T., Prasilova, P. & Pankova, K. (2004) Relationships among vernalization, shoot apex development and frost tolerance in wheat. *Annals of Botany*, 94, 413-418.

Pugsley, A. (1971) A genetic analysis of spring–winter habit of growth in wheat. *Australian Journal of Agricultural Research*, 22, 21-31.

Raman, H., Platz, G. J., Chalmers, K. J., Raman, R., Read, B. J., Barr, A. R. & Moody, D. B. (2003) Mapping of genomic regions associated with net form of net blotch resistance in barley. *Australian Journal of Agricultural Research*, 54, 1359-1367.

Reddy, L., Allan, R. E. & Campbell, K. A. G. (2006) Evaluation of cold hardiness in two sets of near-isogenic lines of wheat (*Triticum aestivum*) with polymorphic vernalization alleles. *Plant Breeding*, 125, 448-456.

Ribaut, J. M., Jiang, C. & Hoisington, D. (2002) Simulation experiments on efficiencies of gene introgression by backcrossing. *Crop Science*, 42, 557-565.

Saisho, D., Kawasaki, S., Sato, K. & Takeda, K. (2002) Construction of a BAC library from Japanese malting barley Haruna Nijo. *Plant and Animal Genome*. San Diego.

Sasaki, T. (2005) The complete rice genome sequence and its application to breeding and genetics. *Rice is life: scientific perspectives for the 21st century*. Proceedings of the World Rice Research Conference held in Tsukuba, Japan, 4-7 November 2004, 66-69.

- Saulescu, N. N. & Braun, H. J. (2001) Cold Tolerance. IN Reynolds, M. P., Ortiz-Monasterio, J. I. & McNab, A. (Eds.) *Application of Physiology in Wheat Breeding*.
- Scherrer, B., Isidore, E., Klein, P., Kim, J. S., Bellec, A., Chalhoub, B., Keller, B. & Feuillet, C. (2005) Large intraspecific haplotype variability at the *Rph7* locus results from rapid and recent divergence in the barley genome. *Plant Cell*, 17, 361-374.
- Shimada, S., Ogawa, T., Kitagawa, S., Suzuki, T., Ikari, C., Shitsukawa, N., Abe, T., Kawahigashi, H., Kikuchi, R., Handa, H. & Murai, K. (2009) A genetic network of flowering-time genes in wheat leaves, in which an *APETALA1/FRUITFULL*-like gene, *VRN1*, is upstream of *FLOWERING LOCUS T*. *Plant Journal*, 58, 668-681.
- Single, W. V. (1988) Resistance to frost injury during stem elongation and early heading. *Frost Injury of Wheat - A workshop of the wheat research council*, 7-17.
- Slafer, G. A. & Rawson, H. M. (1994) Sensitivity of wheat phasic development to major environmental-factors - a reexamination of some assumptions made by physiologists and modelers. *Australian Journal of Plant Physiology*, 21, 393-426.
- Sleper, D. A. & Poehlman, J. M. (2006) *Breeding field crops*, Wiley-Blackwell.
- Snape, J. W., Sarma, R., Quarrie, S. A., Fish, L., Galiba, G. & Sutka, J. (2001) Mapping genes for flowering time and frost tolerance in cereals using precise genetic stocks. *Euphytica*, 120, 309-315.
- Sorrells, M. E., La Rota, M., Bermudez-Kandianis, C. E., Greene, R. A., Kantety, R., Munkvold, J. D., Miftahudin, Mahmoud, A., Ma, X. F., Gustafson, P. J., Qi, L. L. L., Echaliier, B., Gill, B. S., Matthews, D. E., Lazo, G. R., Chao, S. M., Anderson, O. D.,

Edwards, H., Linkiewicz, A. M., Dubcovsky, J., Akhunov, E. D., Dvorak, J., Zhang, D. S., Nguyen, H. T., Peng, J. H., Lapitan, N. L. V., Gonzalez-Hernandez, J. L., Anderson, J. A., Hossain, K., Kalavacharla, V., Kianian, S. F., Choi, D. W., Close, T. J., Dilbirligi, M., Gill, K. S., Steber, C., Walker-Simmons, M. K., Mcguire, P. E. & Qualset, C. O. (2003) Comparative DNA sequence analysis of wheat and rice genomes. *Genome Research*, 13, 1818-1827.

Stam, P. & Zeven, A. C. (1981) The theoretical proportion of the donor genome in near-isogenic lines of self-fertilizers bred by backcrossing. *Euphytica*, 30, 227-238.

Stein, N., Prasad, M., Scholz, U., Thiel, T., Zhang, H. N., Wolf, M., Kota, R., Varshney, R. K., Perovic, D., Grosse, I. & Graner, A. (2007) A 1,000-loci transcript map of the barley genome: new anchoring points for integrative grass genomics. *Theoretical and Applied Genetics*, 114, 823-839.

Sutka, J. (1994) Genetic control of frost tolerance in wheat (*Triticum aestivum* L.). *Euphytica*, 77, 277-282.

Sutka, J. (2001) Genes for frost resistance in wheat. *Euphytica*, 119, 167-172.

Szucs, P., Skinner, J. S., Karsai, I., Cuesta-Marcos, A., Haggard, K. G., Corey, A. E., Chen, T. H. H. & Hayes, P. M. (2007) Validation of the *VRN-H2/VRN-H1* epistatic model in barley reveals that intron length variation in *VRN-H1* may account for a continuum of vernalization sensitivity. *Molecular Genetics and Genomics*, 277, 249-261.

- Takahashi, R. & Yasuda, S. (1971) Genetics of earliness and growth habit in barley. IN Nilan, R. A. (Ed.) 2nd International Barley Genetics Symposium. Washington State University Press, Washington.
- Thomashow, M. F. (1990) Molecular genetics of cold acclimation in higher plants. *Advances In Genetics*, 28, 99-131.
- Toth, B., Galiba, G., Feher, E., Sutka, J. & Snape, J. W. (2003) Mapping genes affecting flowering time and frost resistance on chromosome 5B of wheat. *Theoretical and Applied Genetics*, 107, 509-514.
- Tranquilli, G. & Dubcovsky, J. (2000) Epistatic interaction between vernalization genes *Vrn-A(m)1* and *Vrn-A(m)2* in diploid wheat. *Journal of Heredity*, 91, 304-306.
- Trevaskis, B., Hemming, M. N., Dennis, E. S. & Peacock, W. J. (2007) The molecular basis of vernalization-induced flowering in cereals. *Trends in Plant Science*, 12, 352-357.
- Trevaskis, B., Hemming, M. N., Peacock, W. J. & Dennis, E. S. (2006) *HvVRN2* responds to daylength, whereas *HvVRN1* is regulated by vernalization and developmental status. *Plant Physiology*, 140, 1397-1405.
- Vagujfalvi, A., Galiba, G., Cattivelli, L. & Dubcovsky, J. (2003) The cold-regulated transcriptional activator *Cbf3* is linked to the frost-tolerance locus *Fr-A2* on wheat chromosome 5A. *Molecular Genetics and Genomics*, 269, 60-67.
- Wheeler, R. (2007) Barley variety sowing guide 2008. SARDI cereal crop sowing guides 2008.

- White, C. (2000) Cereals : frost identification : the back pocket guide. Department of Agriculture and Food, Western Australia.
- Williams, J., Hook, R. & Hamblin, A. (2002) Agro-ecological zones of Australia: methodologies for their derivation and key issues in resource management. Canberra, CSIRO Land and Water.
- Williams, K. J. (2003) The molecular genetics of disease resistance in barley. Australian Journal of Agricultural Research, 54, 1065-1079.
- Yan, L., Helguera, M., Kato, K., Fukuyama, S., Sherman, J. & Dubcovsky, J. (2004a) Allelic variation at the *VRN-1* promoter region in polyploid wheat. Theoretical and Applied Genetics, 109, 1677-1686.
- Yan, L., Loukoianov, A., Tranquilli, G., Helguera, M., Fahima, T. & Dubcovsky, J. (2003) Positional cloning of the wheat vernalization gene *VRN1*. Proceedings of the National Academy of Sciences of the United States of America, 100, 6263-6268.
- Yan, L. L., Loukoianov, A., Blechl, A., Tranquilli, G., Ramakrishna, W., Sanmiguel, P., Bennetzen, J. L., Echenique, V. & Dubcovsky, J. (2004b) The wheat *VRN2* gene is a flowering repressor down-regulated by vernalization. Science, 303, 1640-1644.
- Yu, Y., Tomkins, J. P., Waugh, R., Frisch, D. A., Kudrna, D., Kleinhofs, A., Brueggeman, R. S., Muehlbauer, G. J., Wise, R. P. & Wing, R. A. (2000) A bacterial artificial chromosome library for barley (*Hordeum vulgare* L.) and the identification of clones containing putative resistance genes. Theoretical and Applied Genetics, 101, 1093-1099.

Zadoks, J. C., Chang, T. T. & Konzak, C. F. (1974) A decimal code for the growth stages in cereals. *Weed Research*, 14, 415-421.